
## State Highlights 2002


SMOKING-ATTRIBUTABLE DEATHS, 1999		SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
Overall Men	7,412 4,933	State Lung Cancer All States	90.2/100,000	Overall Projected Death	111,273
Women Death Rate	2,479 326.6/100,000	State CHD* All States	60.6/100,000 59.7/100,000	Rate Projected All	9,905/100,000
All States	295.5/100,000	State COPD† All States	73.6/100,000 59.7/100,000	States	8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

#### **Adult Cigarette Use, 2000**



#### Environmental Tobacco Smoke, 1998-99




#### **Youth Tobacco Use**

GRADES 6–8			Grades 9–12	
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
AL†	19.1%	26.5%	30.2%	37.6%
Boys†	22.1%	32.0%	32.5%	44.3%
Girls†	15.7%	20.4%	27.8%	30.7%

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $^{\dagger}$ Alabama Youth Tobacco Survey, 2000.

#### **Disparities Among Adult Population Groups, 2000\***





Alabama is investing 8% of CDC's Best Practices lower estimated recommended funding and 3% of the upper estimated recommended funding. Alabama spent \$269 per capita on smoking-attributable direct medical expenditures. In 1998, about 9% (\$186,000,000 or \$352.36 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

#### **Smoking-Attributable Direct Medical Expenditures, 1998**

#### Ambulatory \$433,000,000 Hospital \$300,000,000 Nursing Home \$241,000,000 Prescription Drugs \$110,000,000 Other \$86,000,000 \$1,170,000,000 Annual Total Annual Per Capita \$269

#### Smoking-Attributable **Productivity Costs, 1999**

Annual Total Annual Per Capita	\$1,777,000,000 \$407

#### State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$96,961,493 \$70,665,000 Gross cigarette tax revenue collected in 2000 Cigarette tax per pack was \$0.165 in 2001

Cigarette sales were 96.2 packs per capita in 2000

Funding Source	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)*	\$600,000	10/01–9/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$600,000	
Federal—CDC Office on Smoking and Health	\$1,340,047	6/01–5/02
Federal—SAMHSA	\$78,720	10/01-9/02
Non-Government Source—American Legacy Foundation	\$208,156	10/00–9/01
Non-Government Source—RWJF/AMA	\$0	
Subtotal: Federal/National Sources	\$1,626,923	
FY02 Total Investment in Tobacco Control		\$2,226,923
CDC Best Practices Recommended Annual Total (Lower )	\$26,740,000	
CDC Best Practices Recommended Annual Total (Upper 1	\$71,235,000	
FY02 Per Capita Investment in Tobacco Control		\$0.50


<sup>\*</sup>An additional \$2,000,000 was appropriated in FY02 to the Department of Education for drug, alcohol, or tobacco education and prevention programs.




SMOKING-ATTRIBUTABLE DEATHS, 1999	SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
Overall       448         Men       264         Women       184         Death Rate       289.6/100,000         All States       295.5/100,000	State Lung Cancer All States State CHD* All States State COPD† All States	88.5/100,000 90.2/100,000 47.1/100,000 59.7/100,000 86.4/100,000 59.7/100,000	Overall Projected Deatl Rate Projected All States	19,516 1 10,233/100,000 8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

#### **Adult Cigarette Use, 2000**



#### **Environmental Tobacco Smoke, 1998-99**




#### **Youth Tobacco Use**

GRADES 6–8			GRADES 9–12	
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
AK <sup>†</sup>	n/a	n/a	36.5%	n/a
Boys <sup>†</sup>	n/a	n/a	31.1%	n/a
Girls <sup>†</sup>	n/a	n/a	35.8%	n/a

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $^{\dagger}$ Alaska Youth Risk Behavior Survey, 1995.

#### Disparities Among Adult Population Groups, 2000\*





Alaska is investing 61% of CDC's *Best Practices* lower estimated recommended funding and 30% of the upper estimated recommended funding. Alaska spent \$215 per capita on smoking-attributable direct medical expenditures. In 1998, about 17% (\$60,000,000 or \$804.31 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

### Smoking-Attributable Direct Medical Expenditures, 1998

# Ambulatory \$54,000,000 Hospital \$40,000,000 Nursing Home \$9,000,000 Prescription Drugs \$9,000,000 Other \$20,000,000 Annual Total \$132,000,000 Annual Per Capita \$215

## Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$129,000,000 \$208

#### State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$21,176,139 Gross cigarette tax revenue collected in 2000 \$43,092,000

Cigarette tax per pack was \$1.00 in 2001


Cigarette sales were 66.0 packs per capita in 2000

FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$3,000,000	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$270,000	7/01–6/02
Subtotal: State Appropriation	\$3,270,000	
Federal—CDC Office on Smoking and Health	\$1,145,052	6/01–5/02
Federal—SAMHSA	\$50,000	10/01–9/02
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$434,752	3/01–2/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		\$4,899,804
CDC Best Practices Recommended Annual Total (Lower F	\$8,088,000	
CDC Best Practices Recommended Annual Total (Upper E	\$16,512,000	
FY02 Per Capita Investment in Tobacco Control	\$7.72	


SMOKING-ATTRIBUTABLE DEATHS, 1999		SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
Overall Men Women Death Rate	6,890 4,382 2,508 283.5/100,000	State Lung Cancer All States State CHD* All States	81.2/100,000 90.2/100,000 58.5/100,000 59.7/100,000	Overall Projected Death Rate Projected All	77,724 5,686/100,000
All States	295.5/100,000	State COPD† All States	76.7/100,000 59.7/100,000	Štates	8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

#### Adult Cigarette Use, 2000



#### **Environmental Tobacco Smoke, 1998–99**




#### **Youth Tobacco Use**

GRADES 6–8			Grades 9–12	
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
AZ <sup>†</sup>	11.4%	17.1%	n/a	n/a
Boys <sup>†</sup>	11.5%	18.0%	n/a	n/a
Girls <sup>†</sup>	11.3%	16.1%	n/a	n/a

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $^{\dagger}$ Arizona Youth Tobacco Survey, 2000.

#### **Disparities Among Adult Population Groups, 2000\***





As of November 30, 2001, the state legislature had not completed its FY2002 budget bill; therefore, investment in tobacco control figures were not available for this report. Arizona spent \$215 per capita on smoking-attributable direct medical expenditures. In 1998, about 14% (\$247,000,000 or \$485.57 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

## Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory	\$435,000,000
Hospital	\$233,000,000
Nursing Home	\$175,000,000
Prescription Drugs	\$94,000,000
Other	\$68,000,000
Annual Total	\$1,005,000,000
Annual Per Capita	\$215

## Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$1,309,000,000 \$274

#### State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$87,669,615 Gross cigarette tax revenue collected in 2000 \$162,795,000

Cigarette tax per pack was \$0.58 in 2001


Cigarette sales were 58.5 packs per capita in 2000

FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE			
State Appropriation—Settlement (Tobacco Only)	not available				
State Appropriation—Excise Tax Revenue	not available				
State Appropriation—Other	not available				
Subtotal: State Appropriation	not available				
Federal—CDC Office on Smoking and Health	\$256,630	6/01–5/02			
Federal—SAMHSA	\$0				
Non-Government Source—American Legacy Foundation	\$91,500	10/00–9/01			
Non-Government Source—RWJF/AMA	\$300,000	6/01–5/02			
Subtotal: Federal/National Sources					
FY02 Total Investment in Tobacco Control	FY02 Total Investment in Tobacco Control				
CDC Best Practices Recommended Annual Total (Lower F	\$27,788,000				
CDC Best Practices Recommended Annual Total (Upper F	\$71,102,000				
FY02 Per Capita Investment in Tobacco Control	not available				




	IBUTABLE DEATHS, 1999	SMOKING-ATTRIBUTABLE DEATHS, 1999 TO DIE FROM SMOKE DISEASE-SPECIFIC 1999–2000		A SMOKING,	
Overall Men Women	4,901 3,113 1,788	State Lung Cancer All States State CHD*	113.2/100,000 90.2/100,000 70.2/100,000	Overall Projected Death Rate	66,957 9,841/100,000
Death Rate All States	343.3/100,000 295.5/100,000	All States State COPD† All States	59.7/100,000 71.8/100,000 59.7/100,000	Projected All States	8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.



#### **Environmental Tobacco Smoke, 1998-99**




#### **Youth Tobacco Use**

GRADES 6–8		GRADE	Current Cigarette Current Any	
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
AR†	15.8%	22.4%	35.8%	43.8%
Boys†	15.7%	25.4%	37.5%	49.9%
Girls†	15.7%	19.1%	33.7%	37.0%

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $^{\dagger}$ Arkansas Youth Tobacco Survey, 2000.

#### **Disparities Among Adult Population Groups, 2000\***





Arkansas is investing 79% of CDC's *Best Practices* lower estimated recommended funding and 30% of the upper estimated recommended funding. Arkansas spent \$250 per capita on smoking-attributable direct medical expenditures. In 1998, about 14% (\$189,000,000 or \$445.13 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

## Smoking-Attributable Direct Medical Expenditures, 1998

## Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$1,122,000,000 \$440

#### State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$51,364,510 Gross cigarette tax revenue collected in 2000 \$82,409,000

Cigarette tax per pack was \$0.315 in 2001


Cigarette sales were 99.4 packs per capita in 2000

FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$11,931,942	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$376,256	7/01–6/02
Subtotal: State Appropriation	\$12,308,198	
Federal—CDC Office on Smoking and Health	\$1,275,017	6/01–5/02
Federal—SAMHSA	\$153,346	10/01–9/02
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$381,867	6/01–5/02
Subtotal: Federal/National Sources	\$1,810,230	
FY02 Total Investment in Tobacco Control		\$14,118,428
CDC Best Practices Recommended Annual Total (Lower E	\$17,906,000	
CDC Best Practices Recommended Annual Total (Upper E	\$46,445,000	
FY02 Per Capita Investment in Tobacco Control	\$5.30	




SMOKING-ATTRIBUTABLE DEA	<sup>THS</sup> , 1	IBUTABLE DEATHS, 1999 E-Specific	EATHS, YOUTH PROJECTED TO DIE FROM SMOKIN 1999–2000	
Men 22		74.1/100,000 90.2/100,000 60.8/100,000 59.7/100,000 68.8/100,000 59.7/100,000	Overall Projected Death Rate Projected All States	637,270 6,890/100,000 8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.



#### **Environmental Tobacco Smoke, 1998-99**




#### **Youth Tobacco Use**

GRADES 6–8		GRADES 9–12		
Current Ci		Current Any	Current Cigarette	Current Any
Smoki		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
CA <sup>†</sup>	6.7%	10.0%	21.6%	27.8%
Boys <sup>†</sup>	7.0%	10.6%	22.7%	31.0%
Girls <sup>†</sup>	6.4%	9.4%	20.4%	24.3%

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $^{\dagger}$ California Youth Tobacco Survey, 2000.

#### **Disparities Among Adult Population Groups, 2000\***





California is investing 83% of CDC's *Best Practices* lower estimated recommended funding and 31% of the upper estimated recommended funding. California spent \$218 per capita on smoking-attributable direct medical expenditures. In 1998, about 16% (\$2,310,000,000 or \$326.14 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

## Smoking-Attributable Direct Medical Expenditures, 1998

# Ambulatory \$3,703,000,000 Hospital \$1,363,000,000 Nursing Home \$1,140,000,000 Prescription Drugs \$507,000,000 Other \$424,000,000 Annual Total \$7,137,000,000 Annual Per Capita \$218

## Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$7,515,000,000 \$227

#### State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$759,225,833 Gross cigarette tax revenue collected in 2000 \$1,176,859,000

Cigarette tax per pack was \$0.87 in 2001

Cigarette sales were 41.6 packs per capita in 2000


Funding Source	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)*	\$20,000,000	7/01–6/02
State Appropriation—Excise Tax Revenue	\$114,536,000	7/01–6/02
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$134,536,000	
Federal—CDC Office on Smoking and Health	\$335,610	6/01–5/02
Federal—SAMHSA	\$2,000,000	10/01–9/02
Non-Government Source—American Legacy Foundation	\$645,679	10/00–9/01
Non-Government Source—RWJF/AMA	\$299,176	6/01–5/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		\$137,816,465
CDC Best Practices Recommended Annual Total (Lower 1	\$165,098,000	
CDC Best Practices Recommended Annual Total (Upper 1	\$442,403,000	
FY02 Per Capita Investment in Tobacco Control	\$4.02	

<sup>\*</sup>This amount represents one state's portion of the tobacco settlement revenue.


SMOKING-ATTRIBUTABLE DEATHS, 1999	1999	SMOKING-ATTRIBUTABLE DEATHS, 1999 TO DIE FROM SM DISEASE-SPECIFIC 1999–200		M SMOKING,
Overall         4,276           Men         2,537           Women         1,739           Death Rate         248.1/100,000           All States         295.5/100,000	State Lung Cancer All States State CHD* All States State COPD† All States	62.0/100,000 90.2/100,000 41.5/100,000 59.7/100,000 85.2/100,000 59.7/100,000	Overall Projected Death Rate Projected All States	92,246 8,380/100,000 8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

#### Adult Cigarette Use, 2000



#### **Environmental Tobacco Smoke, 1998–99**




#### **Youth Tobacco Use**

GRADES 6–8		GRADE	s <b>9–12</b>	
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
CO <sup>†</sup>	8.8%	13.6%	25.3%	34.4%
Boys <sup>†</sup>	9.3%	15.9%	23.9%	37.1%
Girls <sup>†</sup>	8.4%	11.1%	26.9%	31.6%

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $^{\dagger}$ Colorado Youth Tobacco Survey, 2000.

#### **Disparities Among Adult Population Groups, 2000\***





Colorado is investing 61% of CDC's *Best Practices* lower estimated recommended funding and 24% of the upper estimated recommended funding. Colorado spent \$259 per capita on smoking-attributable direct medical expenditures. In 1998, about 17% (\$249,000,000 or \$722.84 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

### Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory Hospital Nursing Home Prescription Drugs Other Annual Total Annual Per Capita	\$419,000,000 \$239,000,000 \$213,000,000 \$73,000,000 \$82,000,000 \$1,026,000,000 \$259
Annual Per Capita	\$259

## Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$856,000,000 \$211

#### State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$85,026,250 Gross cigarette tax revenue collected in 2000 \$60,818,000

Cigarette tax per pack was \$0.20 in 2001

Cigarette sales were 73.0 packs per capita in 2000

FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)*	\$12,891,329	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$12,891,329	
Federal—CDC Office on Smoking and Health	\$1,350,347	6/01–5/02
Federal—SAMHSA	\$99,940	10/01-9/02
Non-Government Source—American Legacy Foundation†	10/00–9/01	
Non-Government Source—RWJF/AMA	\$383,176	3/01–2/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control	\$14,924,792	
CDC Best Practices Recommended Annual Total (Lower E	\$24,546,000	
CDC Best Practices Recommended Annual Total (Upper E	\$63,255,000	
FY02 Per Capita Investment in Tobacco Control	\$3.39	


<sup>\*</sup>An additional \$6,875,375 was appropriated to the Office of the President at the University of Colorado for a tobacco-related research grant program. †On October 1, 2001, Colorado was awarded an additional grant from the American Legacy Foundation.




Smoking-Attributable Deaths, 1999		SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
Overall Men Women Death Rate 255.9/10 All States 295.5/10		State Lung Cancer All States State CHD* All States State COPD† All States	78.6/100,000 90.2/100,000 54.7/100,000 59.7/100,000 55.6/100,000 59.7/100,000	Overall Projected Death Rate Projected All States	83,859 9,963/100,000 8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

#### **Adult Cigarette Use, 2000**



#### **Environmental Tobacco Smoke, 1998–99**




#### **Youth Tobacco Use**

GRADES 6–8		Grades 9–12		
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
CT <sup>†</sup>	9.8%	13.1%	25.6%	32.4%
Boys <sup>†</sup>	9.7%	14.1%	24.9%	35.3%
Girls <sup>†</sup>	9.8%	11.9%	26.0%	29.2%

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $^{\dagger}$ Connecticut Youth Tobacco Survey, 2000.

#### Disparities Among Adult Population Groups, 2000\*





Connecticut is investing 12% of CDC's *Best Practices* lower estimated recommended funding and 5% of the upper estimated recommended funding. Connecticut spent \$389 per capita on smoking-attributable direct medical expenditures. In 1998, about 13% (\$336,000,000 or \$880.85 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

### Smoking-Attributable Direct Medical Expenditures, 1998

# Ambulatory \$383,000,000 Hospital \$211,000,000 Nursing Home \$496,000,000 Prescription Drugs \$95,000,000 Other \$88,000,000 Annual Total \$1,273,000,000 Annual Per Capita \$389

## Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$870,000,000 \$265

#### State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$110,430,429 Gross cigarette tax revenue collected in 2000 \$117,215,000

Cigarette tax per pack was \$0.50 in 2001

Cigarette sales were 71.4 packs per capita in 2000


FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)*	\$1,046,550	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$200,000	7/01–6/02
Subtotal: State Appropriation	\$1,246,550	
Federal—CDC Office on Smoking and Health	\$1,010,252	6/01–5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$316,727	3/01–6/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control	\$2,573,529	
CDC Best Practices Recommended Annual Total (Lower	\$21,240,000	
CDC Best Practices Recommended Annual Total (Upper	\$53,895,000	
FY02 Per Capita Investment in Tobacco Control	<b>\$0.75</b>	

 $<sup>*</sup> An additional \$13,\!500,\!000 \ was \ appropriated \ to \ the \ Tobacco \ \& \ Health \ Trust \ Fund \ for \ tobacco \ control \ and \ other \ health \ programs.$ 


	IBUTABLE DEATHS, 1999	SMOKING-ATTRIBU 1999 DISEASE-SI	<b>)</b>	TO DIE FRO	ROJECTED M SMOKING, –2000
Overall Men Women Death Rate	1,193 708 485 313.9/100,000	State Lung Cancer All States State CHD* All States	113.1/100,000 90.2/100,000 67.5/100,000 59.7/100,000	Overall Projected Death Rate Projected All	19,584 10,064/100,000
All States	295.5/100,000	State COPD† All States	59.7/100,000 66.6/100,000 59.7/100,000	States	8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

#### Adult Cigarette Use, 2000



#### Environmental Tobacco Smoke, 1998-99




#### **Youth Tobacco Use**

Grades 6–8		GRADES 9–12		
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
DE†	15.2%	17.8%	27.1%	31.2%
Boys†	14.8%	18.9%	28.0%	34.6%
Girls†	15.6%	16.6%	26.1%	27.8%

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $^{\dagger}$ Delaware Youth Tobacco Survey, 2000.

#### **Disparities Among Adult Population Groups, 2000\***





Delaware is investing 72% of CDC's *Best Practices* lower estimated recommended funding and 34% of the upper estimated recommended funding. Delaware spent \$298 per capita on smoking-attributable direct medical expenditures. In 1998, about 16% (\$62,000,000 or \$609.71 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

#### Smoking-Attributable Direct Medical Expenditures, 1998

## Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$258,000,000 \$342

#### State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$23,523,288 Gross cigarette tax revenue collected in 2000 \$25,957,000

Cigarette tax per pack was \$0.24 in 2001

Cigarette sales were 140.7 packs per capita in 2000


Funding Source	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$5,454,900	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$5,454,900	
Federal—CDC Office on Smoking and Health	\$782,761	6/01–5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$0	
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		\$6,237,661
CDC Best Practices Recommended Annual Total (Lower E	\$8,631,000	
CDC Best Practices Recommended Annual Total (Upper E	\$18,464,000	
FY02 Per Capita Investment in Tobacco Control	\$7.85	




SMOKING-ATTRIBUTABLE DEATHS, 1999	SMOKING-ATTRIBUT 1999 DISEASE-SI	)	HS, YOUTH PROJECTED TO DIE FROM SMOKIN 1999–2000	
Overall         721           Men         456           Women         265           Death Rate         250.4/100,000           All States         295.5/100,000	State CHD*	80.0/100,000 90.2/100,000 50.4/100,000 59.7/100,000 38.7/100,000 59.7/100,000	Overall Projected Death Rate Projected All States	7,786 6,771/100,000 8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

#### **Adult Cigarette Use, 2000**



#### **Environmental Tobacco Smoke, 1998-99**




#### **Youth Tobacco Use**

GRADES 6–8		Grades 9–12		
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
DC <sup>†</sup>	9.4%	14.2%	14.7%	21.0%
Boys <sup>†</sup>	8.7%	15.2%	16.9%	24.9%
Girls <sup>†</sup>	10.1%	13.4%	12.3%	16.5%

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000; †District of Columbia Youth Tobacco Survey, 2000.

#### Disparities Among Adult Population Groups, 2000\*





The District of Columbia is investing 8% of CDC's *Best Practices* lower estimated recommended funding and 4% of the upper estimated recommended funding. The District of Columbia spent \$364 per capita on smoking-attributable direct medical expenditures. In 1998, about 9% (\$61,000,000 or \$367.58 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

## Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory	\$58,000,000
Hospital	\$78,000,000
Nursing Home	\$37,000,000
Prescription Drugs	\$8,000,000
Other	\$9,000,000
Annual Total	\$190,000,000
Annual Per Capita	\$364

## Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$212,000,000 \$408

#### State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001

not available \$17,050,000

Gross cigarette tax revenue collected in 2000

Cigarette tax per pack was \$0.65 in 2001


Cigarette sales were 50.2 packs per capita in 2000

FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE		
State Appropriation—Settlement (Tobacco Only)	\$0			
State Appropriation—Excise Tax Revenue	\$0			
State Appropriation—Other	\$0			
Subtotal: State Appropriation	\$0			
Federal—CDC Office on Smoking and Health	\$448,157	6/01–5/02		
Federal—SAMHSA	\$36,292	10/01-9/02		
Non-Government Source—American Legacy Foundation	\$100,000	10/00-9/02		
Non-Government Source—RWJF/AMA	\$0			
Subtotal: Federal/National Sources	\$584,449			
FY02 Total Investment in Tobacco Control	\$584,449			
CDC Best Practices Recommended Annual Total (Lower F	\$7,479,000			
CDC Best Practices Recommended Annual Total (Upper E	\$14,571,000			
FY02 Per Capita Investment in Tobacco Control				




SMOKING-ATTRIBUTABLE DEATH	SMOKING-ATTRIBU 199 DISEASE-S	9	Youth Pr to Die from 1999–	SMOKING,
Overall       28,7         Men       17,4         Women       11,2         Death Rate       283.1/100,0         All States       295.5/100,0	All States State CHD* All States	91.9/100,000 90.2/100,000 64.2/100,000 59.7/100,000 65.6/100,000 59.7/100,000	Overall Projected Death Rate Projected All States	306,724 8,412/100,000 8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.



#### **Environmental Tobacco Smoke, 1998–99**




#### **Youth Tobacco Use**

Grades 6–8		GRADE	s 9–12	
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
FL <sup>†</sup>	9.8%	13.5%	19.0%	25.8%
Boys <sup>†</sup>	10.0%	15.6%	19.1%	29.6%
Girls <sup>†</sup>	9.5%	11.2%	19.0%	22.0%

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $\dagger$ Florida Youth Tobacco Survey, 2000.

#### **Disparities Among Adult Population Groups, 2000\***





Florida is investing 39% of CDC's Best Practices lower estimated recommended funding and 14% of the upper estimated recommended funding. Florida spent \$331 per capita on smoking-attributable direct medical expenditures. In 1998, about 16% (\$976,000,000 or \$512.30 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

## Smoking-Attributable Direct Medical Expenditures, 1998

# Ambulatory \$1,916,000,000 Hospital \$948,000,000 Nursing Home \$1,194,000,000 Prescription Drugs \$478,000,000 Other \$398,000,000 Annual Total \$4,934,000,000 Annual Per Capita \$331

## Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$5,441,000,000 \$360

#### State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 Gross cigarette tax revenue collected in 2000	\$731,300,000 \$426,269,000	
Cigarette tax per pack was \$0.339 in 2001 Cigarette sales were 82.5 packs per capita in 2000	,,,,	


Funding Source	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$29,803,000	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$29,803,000	
Federal—CDC Office on Smoking and Health	\$750,000	6/01–5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$50,004	10/00–9/01
Non-Government Source—RWJF/AMA	\$0	
Subtotal: Federal/National Sources	\$800,004	
FY02 Total Investment in Tobacco Control		\$30,603,004
CDC Best Practices Recommended Annual Total (Lower E	stimate)	\$78,383,000
CDC Best Practices Recommended Annual Total (Upper E	\$221,260,000	
FY02 Per Capita Investment in Tobacco Control	\$1.88	




SMOKING-ATTRIBUTABLE . 1999	DEATHS,	SMOKING-ATTRIBUT 1999 DISEASE-SP		Youth Pr to Die from 1999–	SMOKING,
	10,728 6,911 3,817 (100,000 (100,000	State Lung Cancer All States State CHD* All States State COPD† All States	99.3/100,000 90.2/100,000 62.5/100,000 59.7/100,000 76.2/100,000 59.7/100,000	Overall Projected Death Rate Projected All States	187,782 8,657/100,000 8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

#### Adult Cigarette Use, 2000



#### **Environmental Tobacco Smoke, 1998-99**




#### **Youth Tobacco Use**

Grades 6–8		GRADE	s 9–12	
Current C		Current Any	Current Cigarette	Current Any
Smol		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
GA	13.8% <sup>†</sup>	18.8% <sup>†</sup>	24.3% <sup>††</sup>	n/a
Boys	13.9% <sup>†</sup>	20.6% <sup>†</sup>	24.7% <sup>††</sup>	n/a
Girls	13.6% <sup>†</sup>	16.5% <sup>†</sup>	24.0% <sup>††</sup>	n/a

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $^{\dagger}$ Georgia Youth Tobacco Survey, 1999;  $^{\dagger}$ Georgia Youth Risk Behavior Survey, 1993.

#### **Disparities Among Adult Population Groups, 2000\***





Georgia is investing 56% of CDC's *Best Practices* lower estimated recommended funding and 21% of the upper estimated recommended funding. Georgia spent \$230 per capita on smoking-attributable direct medical expenditures. In 1998, about 13% (\$419,000,000 or \$343.28 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

## Smoking-Attributable Direct Medical Expenditures, 1998

# Ambulatory \$742,000,000 Hospital \$421,000,000 Nursing Home \$311,000,000 Prescription Drugs \$149,000,000 Other \$135,000,000 Annual Total \$1,758,000,000 Annual Per Capita \$230

## Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$2,734,000,000 \$351

#### State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$152,235,170 Gross cigarette tax revenue collected in 2000 \$67,064,000

Cigarette tax per pack was \$0.12 in 2001


Cigarette sales were 70.9 packs per capita in 2000

FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$20,765,890	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$20,765,890	
Federal—CDC Office on Smoking and Health	\$1,608,025	6/01–5/02
Federal—SAMHSA	\$446,371	10/01–9/02
Non-Government Source—American Legacy Foundation	\$767,286	10/00–9/01
Non-Government Source—RWJF/AMA	\$316,022	6/01–5/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		\$23,903,594
CDC Best Practices Recommended Annual Total (Lower E	\$42,591,000	
CDC Best Practices Recommended Annual Total (Upper E	\$114,314,000	
FY02 Per Capita Investment in Tobacco Control	\$2.86	




SMOKING-ATTRIBUTABLE DEATHS, 1999		SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
Overall Men Women Death Rate All States	1,095 770 325 174.3/100,000	State Lung Cancer All States State CHD* All States State COPD†	53.7/100,000 90.2/100,000 35.2/100,000 59.7/100,000	Overall Projected Death Rate Projected All States	23,858 8,067/100,000
An States	295.5/100,000	All States	30.1/100,000 59.7/100,000	States	8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.



#### **Environmental Tobacco Smoke, 1998-99**




#### **Youth Tobacco Use**

Grades 6–8			GRADES 9–12		
Current Cigarette Current Any Smoking Tobacco Use		Current Cigarette Smoking	Current Any Tobacco Use		
National* HI <sup>†</sup> Boys <sup>†</sup> Girls <sup>†</sup>	11.0% n/a n/a n/a	15.1% n/a n/a n/a	28.0% 24.5% 23.3% 25.5%	34.5% 27.1% 26.2% 27.7%	

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $\dagger$ Hawaii Youth Tobacco Survey, 2000.

#### **Disparities Among Adult Population Groups, 2000\***





Hawaii is investing 217% of CDC's Best Practices lower estimated recommended funding and 100% of the upper estimated recommended funding. Hawaii spent \$220 per capita on smoking-attributable direct medical expenditures. In 1998, about 14% (\$91,000,000 or \$493.42 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

## Smoking-Attributable Direct Medical Expenditures, 1998

## Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$263,000,000 \$222

#### State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$35,800,091 Gross cigarette tax revenue collected in 2000 \$40,050,000

Cigarette tax per pack was \$1.00 in 2001

Cigarette sales were 31.9 packs per capita in 2000


FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)*	\$22,171,571	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$22,171,571	
Federal—CDC Office on Smoking and Health	\$874,172	6/01–5/02
Federal—SAMHSA	\$68,989	10/01–9/02
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$283,057	6/01–5/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control	\$23,397,789	
CDC Best Practices Recommended Annual Total (Lower E	\$10,778,000	
CDC Best Practices Recommended Annual Total (Upper E	\$23,448,000	
FY02 Per Capita Investment in Tobacco Control	\$19.16	

<sup>\*</sup>The amount reflected above is disbursed to a trust fund. Up to an additional \$9,581,399 was appropriated to the Department of Health for promotion and disease prevention programs, including but not limited to maternal and child health and child development programs, promotion of healthy lifestyles and prevention-oriented public health programs.




SMOKING-ATTRIBUTABLE DEATHS, 1999		Smoking-Attributable Deaths, 1999 Disease-Specific		Youth Projected to Die from Smoking, 1999–2000	
Overall Men	1,505 940	State Lung Cancer All States	68.1/100,000 90.2/100,000	Overall Projected Death	32,127
Women Death Rate	565 254.6/100,000	State CHD* All States	49.5/100,000 59.7/100,000	Rate Projected All	8,706/100,000
All States	295.5/100,000	State COPD† All States	73.7/100,000 59.7/100,000	Štates	8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.



#### **Environmental Tobacco Smoke, 1998-99**




#### **Youth Tobacco Use**

GRADES 6–8		GRADES 9–12		
	Current Cigarette Current Any Current Cigarette Smoking Tobacco Use Smoking		Current Any Tobacco Use	
National* ID Boys Girls	11.0% 9.6% <sup>†</sup> 9.7% <sup>†</sup> 9.4% <sup>†</sup>	15.1% 13.4% <sup>†</sup> 15.4% <sup>†</sup> 11.1% <sup>†</sup>	28.0% 27.3% <sup>††</sup> 29.3% <sup>††</sup> 25.5% <sup>††</sup>	34.5% n/a n/a n/a

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000; †Idaho Youth Tobacco Survey, 2001; ††Idaho Youth Risk Behavior Survey, 1993.

#### **Disparities Among Adult Population Groups, 2000\***





Idaho is investing 24% of CDC's Best Practices lower estimated recommended funding and 11% of the upper estimated recommended funding. Idaho spent \$202 per capita on smoking-attributable direct medical expenditures. In 1998, about 14% (\$65,000,000 or \$527.33 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

#### **Smoking-Attributable Direct Medical Expenditures, 1998**

#### **Smoking-Attributable Productivity Costs, 1999**

Annual Total Annual Per Capita	\$271,000,000 \$217

#### State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$22,531,022 Gross cigarette tax revenue collected in 2000 \$25,250,000

Cigarette tax per pack was \$0.28 in 2001

Cigarette sales were 66.9 packs per capita in 2000


FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)*	\$1,006,500	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$300,000	7/01–6/02
Subtotal: State Appropriation	\$1,306,500	
Federal—CDC Office on Smoking and Health	\$1,113,388	6/01–5/02
Federal—SAMHSA	\$50,000	10/01-9/02
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$200,000	6/01–5/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control	\$2,669,888	
CDC Best Practices Recommended Annual Total (Lower I	\$11,044,000	
CDC Best Practices Recommended Annual Total (Upper I	\$24,087,000	
FY02 Per Capita Investment in Tobacco Control	\$2.02	

<sup>\*</sup>An additional \$50,000 was appropriated to the Buhl School District Substance Abuse Pilot Program to address tobacco and/or substance abuse. An additional \$120,000 was appropriated to the Idaho Supreme Court to address tobacco and/or substance abuse.




SMOKING-ATTRIBUTABLE DEAT	199	SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
Overall       18,         Men       10,         Women       7,         Death Rate       303.1/100,         All States       295.5/100,	47 All States 29 State CHD* 00 All States	r 91.0/100,000 90.2/100,000 69.6/100,000 59.7/100,000 63.7/100,000 59.7/100,000	Overall Projected Death Rate Projected All States	297,384 9,163/100,000 8,830/100,000	

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.



#### **Environmental Tobacco Smoke, 1998-99**




#### **Youth Tobacco Use**

	GRADI	ES 6–8	GRADE	s 9–12
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
IL†	n/a	n/a	35.7%	n/a
Boys†	n/a	n/a	35.8%	n/a
Girls†	n/a	n/a	35.6%	n/a

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $\dagger$ Illinois Youth Risk Behavior Survey, 1995.

#### **Disparities Among Adult Population Groups, 2000\***





Illinois is investing 78% of CDC's Best Practices lower estimated recommended funding and 28% of the upper estimated recommended funding. Illinois spent \$266 per capita on smoking-attributable direct medical expenditures. In 1998, about 18% (\$1,226,000,000 or \$899.06 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

### Smoking-Attributable Direct Medical Expenditures, 1998

# Ambulatory \$1,097,000,000 Hospital \$801,000,000 Nursing Home \$846,000,000 Prescription Drugs \$268,000,000 Other \$194,000,000 Annual Total \$3,206,000,000 Annual Per Capita \$266

## Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$3,909,000,000 \$322

#### State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$288,674,835 Gross cigarette tax revenue collected in 2000 \$485,041,000

Cigarette tax per pack was \$0.58 in 2001

Cigarette sales were 70.0 packs per capita in 2000


FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)*	\$48,900,000	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$48,900,000	
Federal—CDC Office on Smoking and Health	\$1,655,389	6/01–5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$265,960	6/01–5/02
Subtotal: Federal/National Sources	\$1,921,349	
FY02 Total Investment in Tobacco Control		\$50,821,349
CDC Best Practices Recommended Annual Total (Lower E	\$64,909,000	
CDC Best Practices Recommended Annual Total (Upper E	\$179,048,000	
FY02 Per Capita Investment in Tobacco Control		\$4.06

<sup>\*</sup>An additional \$2,250,000 was appropriated for school-based clinics.




Smoking-Attribu	· · · · · · · · · · · · · · · · · · ·	SMOKING-ATTRIBUT 1999 DISEASE-ST	<b>)</b>	to Die fra	Projected Om Smoking, D–2000
Overall Men Women Death Rate All States	10,271 6,146 4,125 341.4/100,000 295.5/100,000	State Lung Cancer All States State CHD* All States State COPD† All States	107.1/100,000 90.2/100,000 71.0/100,000 59.7/100,000 78.1/100,000 59.7/100,000	Overall Projected Death Rate Projected All States	167,604 1 10,646/100,000 8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.



#### **Environmental Tobacco Smoke, 1998-99**




#### **Youth Tobacco Use**

	GRADI	ES <b>6–8</b>	GRADES	s <b>9–12</b>
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
IN <sup>†</sup>	9.8%	15.3%	31.6%	36.9%
Boys <sup>†</sup>	9.3%	16.3%	32.8%	41.1%
Girls <sup>†</sup>	10.4%	14.2%	30.1%	32.3%

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $\dagger$ Indiana Youth Tobacco Survey, 2000.

#### **Disparities Among Adult Population Groups, 2000\***





Indiana is investing 19% of CDC's *Best Practices* lower estimated recommended funding and 7% of the upper estimated recommended funding. Indiana spent \$275 per capita on smoking-attributable direct medical expenditures. In 1998, about 15% (\$380,000,000 or \$625.31 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

## Smoking-Attributable Direct Medical Expenditures, 1998

# Ambulatory \$501,000,000 Hospital \$419,000,000 Nursing Home \$500,000,000 Prescription Drugs \$134,000,000 Other \$73,000,000 Annual Total \$1,627,000,000 Annual Per Capita \$275

## Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$2,164,000,000 \$364

#### State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$106,515,675 Gross cigarette tax revenue collected in 2000 \$117,604,000

Cigarette tax per pack was \$0.155 in 2001

Cigarette sales were 125.5 packs per capita in 2000


FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)*	\$5,000,000	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$5,000,000	
Federal—CDC Office on Smoking and Health	\$1,492,125	6/01–5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$200,000	6/01–5/02
Subtotal: Federal/National Sources	\$1,692,125	
FY02 Total Investment in Tobacco Control		\$6,692,125
CDC Best Practices Recommended Annual Total (Lower E	\$34,784,000	
CDC Best Practices Recommended Annual Total (Upper E	\$95,804,000	
FY02 Per Capita Investment in Tobacco Control		\$1.09

<sup>\*</sup>Funding dedicated to tobacco control from settlement revenue is deposited into a trust fund. In FY01, \$35,000,000 was appropriated to tobacco control. The FY01 appropriation will be rolled over into the FY02 budget.




SMOKING-ATTRIBUTABLE DEATH 1999	1999	Smoking-Attributable Deaths, 1999 Disease-Specific		Projected om Smoking, 9–2000
Overall       4,6         Men       2,9         Women       1,6         Death Rate       265.6/100,0         All States       295.5/100,0	All States State CHD* All States	79.7/100,000 90.2/100,000 57.7/100,000 59.7/100,000 68.6/100,000 59.7/100,000	Overall Projected Deat Rate Projected All States	80,319 h 10,948/100,000 8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.



#### **Environmental Tobacco Smoke, 1998-99**




#### **Youth Tobacco Use**

	GRADI	ES <b>6–8</b>	GRADE	s 9–12
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
IA <sup>†</sup>	11.8%	16.4%	32.7%	39.0%
Boys <sup>†</sup>	11.3%	17.5%	34.2%	45.6%
Girls <sup>†</sup>	12.3%	15.1%	31.1%	31.9%

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $\dagger$ Iowa Youth Tobacco Survey, 2000.

#### **Disparities Among Adult Population Groups, 2000\***





Iowa is investing 58% of CDC's Best Practices lower estimated recommended funding and 23% of the upper estimated recommended funding. Iowa spent \$278 per capita on smoking-attributable direct medical expenditures. In 1998, about 14% (\$235,000,000 or \$745.39 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

## Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory Hospital Nursing Home Prescription Drugs Other Annual Total	\$220,000,000 \$190,000,000 \$264,000,000 \$68,000,000 \$52,000,000 \$794,000,000
Annual Total Annual Per Capita	\$794,000,000 \$278
7 Hillian Fer Capita	Ψ210

## Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$824,000,000 \$287

#### State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$53,940,181 Gross cigarette tax revenue collected in 2000 \$92,817,000

Cigarette tax per pack was \$0.36 in 2001


Cigarette sales were 88.9 packs per capita in 2000

FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$9,420,394	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other		
Subtotal: State Appropriation	\$9,420,394	
Federal—CDC Office on Smoking and Health	\$1,043,669	6/01–5/02
Federal—SAMHSA	\$663,250	10/01-9/02
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA		
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control	\$11,127,313	
CDC Best Practices Recommended Annual Total (Lower F	\$19,347,000	
CDC Best Practices Recommended Annual Total (Upper E	\$48,713,000	
FY02 Per Capita Investment in Tobacco Control	\$3.78	




SMOKING-ATTRIBUTABLE DEATHS 1999	SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
Overall       3,864         Men       2,510         Women       1,354         Death Rate       269.8/100,000         All States       295.5/100,000	All States State CHD* All States	83.6/100,000 90.2/100,000 48.1/100,000 59.7/100,000 70.1/100,000 59.7/100,000	Overall Projected Death Rate Projected All States	57,358 8,045/100,000 8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.



#### **Environmental Tobacco Smoke, 1998-99**




#### **Youth Tobacco Use**

GRADES 6–8		GRADES 9–12		
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
KS†	8.1%	12.0%	26.1%	33.6%
Boys†	8.0%	13.2%	24.7%	37.2%
Girls†	7.9%	10.3%	27.5%	29.7%

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $^{\dagger}$ Kansas Youth Tobacco Survey, 2000.

#### **Disparities Among Adult Population Groups, 2000\***





Kansas is investing 15% of CDC's Best Practices lower estimated recommended funding and 6% of the upper estimated recommended funding. Kansas spent \$274 per capita on smoking-attributable direct medical expenditures. In 1998, about 15% (\$153,000,000 or \$632.56 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

### Smoking-Attributable Direct Medical Expenditures, 1998

## Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$741,000,000 \$279

#### State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$51,707,586 Gross cigarette tax revenue collected in 2000 \$50,887,000

Cigarette tax per pack was \$0.24 in 2001


Cigarette sales were 79.8 packs per capita in 2000

FUNDING SOURCE	FUNDING CYCLE		
State Appropriation—Settlement (Tobacco Only)	\$500,000	7/01–6/02	
State Appropriation—Excise Tax Revenue	State Appropriation—Excise Tax Revenue \$0		
State Appropriation—Other			
Subtotal: State Appropriation	\$500,000		
Federal—CDC Office on Smoking and Health	\$1,229,700	6/01–5/02	
Federal—SAMHSA	\$145,000	10/01–9/02	
Non-Government Source—American Legacy Foundation	\$500,000	10/00–9/01	
Non-Government Source—RWJF/AMA	6/01–5/02		
Subtotal: Federal/National Sources			
FY02 Total Investment in Tobacco Control		\$2,771,608	
CDC Best Practices Recommended Annual Total (Lower E	\$18,052,000		
CDC Best Practices Recommended Annual Total (Upper E	\$44,689,000		
FY02 Per Capita Investment in Tobacco Control	\$1.02		




SMOKING-ATTRIBUTABLE DEATHS, 1999		SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
	7,791 4,836 2,955 100,000 100,000	State Lung Cancer All States State CHD* All States State COPD†	121.4/100,000 90.2/100,000 78.6/100,000 59.7/100,000 91.6/100,000	Overall Projected Death Rate Projected All States	114,995 11,559/100,000 8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.



#### **Environmental Tobacco Smoke, 1998-99**




#### **Youth Tobacco Use**

GRADES 6–8		Grades 9–12		
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
KY†	21.5%	28.3%	37.4%	46.2%
Boys†	21.9%	33.0%	35.6%	50.2%
Girls†	21.1%	23.2%	39.0%	41.8%

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $\dagger$ Kentucky Youth Tobacco Survey, 2000.

#### **Disparities Among Adult Population Groups, 2000\***





Kentucky is investing 20% of CDC's *Best Practices* lower estimated recommended funding and 7% of the upper estimated recommended funding. Kentucky spent \$298 per capita on smoking-attributable direct medical expenditures. In 1998, about 15% (\$380,000,000 or \$589.46 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

# Smoking-Attributable Direct Medical Expenditures, 1998

# Ambulatory \$371,000,000 Hospital \$313,000,000 Nursing Home \$299,000,000 Prescription Drugs \$108,000,000 Other \$80,000,000 Annual Total \$1,171,000,000 Annual Per Capita \$298

# Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$1,841,000,000 \$465

# State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$104,957,133 Gross cigarette tax revenue collected in 2000 \$18,724,000

Cigarette tax per pack was \$0.03 in 2001

Cigarette sales were 156.2 packs per capita in 2000

FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)*	\$3,040,000	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$3,040,000	
Federal—CDC Office on Smoking and Health	\$1,128,413	6/01–5/02
Federal—SAMHSA	\$182,208	10/01-9/02
Non-Government Source—American Legacy Foundation	\$99,261	10/00–9/01
Non-Government Source—RWJF/AMA	3/01–2/02	
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		\$4,912,696
CDC Best Practices Recommended Annual Total (Lower F	\$25,090,000	
CDC Best Practices Recommended Annual Total (Upper F	\$69,895,000	
FY02 Per Capita Investment in Tobacco Control	\$1.21	


<sup>\*</sup>An additional \$2,500,000 was appropriated to the Kentucky Agency for Substance Abuse for smoking cessation and other substance abuse programs.




SMOKING-ATTRIBUTABLE DEATHS, 1999		SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
Overall Men Women	6,421 3,896 2,525	State Lung Cancer All States State CHD*	104.6/100,000 90.2/100,000 65.8/100,000	Overall Projected Death Rate	107,968 8,851/100,000
Death Rate All States	314.1/100,000 295.5/100,000	All States State COPD† All States	59.7/100,000 60.1/100,000 59.7/100,000	Projected All States	8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

# **Adult Cigarette Use, 2000**



# **Environmental Tobacco Smoke, 1998–99**




#### **Youth Tobacco Use**

Grades 6–8			GRADES 9–12		
Current Cigarette Smoking		Current Any Tobacco Use	Current Cigarette Current A Smoking Tobacco U		
National* LA Boys Girls	11.0% 17.1% <sup>†</sup> 18.1% <sup>†</sup> 15.4% <sup>†</sup>	15.1% 26.3% <sup>†</sup> 30.6% <sup>†</sup> 20.8% <sup>†</sup>	28.0% 36.4% <sup>††</sup> 38.2% <sup>††</sup> 34.6% <sup>††</sup>	34.5% n/a n/a n/a	

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000; †Louisiana Youth Tobacco Survey, 2001; ††Louisiana Youth Risk Behavior Survey, 1997.

# **Disparities Among Adult Population Groups, 2000\***





Louisiana is investing 9% of CDC's *Best Practices* lower estimated recommended funding and 3% of the upper estimated recommended funding. Louisiana spent \$264 per capita on smoking-attributable direct medical expenditures. In 1998, about 17% (\$518,000,000 or \$719.08 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

# Smoking-Attributable Direct Medical Expenditures, 1998

# Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$1,662,000,000 \$380

### **State Fiscal Resources Available for Tobacco Control**

Tobacco settlement revenue received in 2001 \$139,997,812 Gross cigarette tax revenue collected in 2000 \$92,299,000

Cigarette tax per pack was \$0.24 in 2001

Cigarette sales were 104.3 packs per capita in 2000


FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$500,000	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$500,000	
Federal—CDC Office on Smoking and Health	\$1,140,677	6/01–5/02
Federal—SAMHSA	\$400,000	10/01-9/02
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	6/01–5/02	
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		\$2,339,851
CDC Best Practices Recommended Annual Total (Lower E	\$27,132,000	
CDC Best Practices Recommended Annual Total (Upper E	\$71,431,000	
FY02 Per Capita Investment in Tobacco Control	\$0.52	




SMOKING-ATTRIBUTABLE DEATHS, 1999		SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
Overall Men Women	2,132 1,216 916	State Lung Cancer All States State CHD*	95.7/100,000 90.2/100,000 56.7/100,000	Overall Projected Deatl Rate	33,292 11,052/100,000
Death Rate All States	305.2/100,000 295.5/100,000	All States State COPD† All States	59.7/100,000 82.5/100,000 59.7/100,000	Projected All States	8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

# Adult Cigarette Use, 2000



#### **Environmental Tobacco Smoke, 1998-99**




#### **Youth Tobacco Use**

Grades 6–8			GRADES 9–12		
Current Cigarette		Current Any	Current Cigarette Current Ar		
Smoking		Tobacco Use	Smoking Tobacco U		
National*	11.0%	15.1%	28.0%	34.5%	
ME†	11.7%	13.7%	28.6%	32.6%	
Boys†	11.3%	14.4%	28.6%	35.2%	
Girls†	12.3%	12.9%	28.5%	29.9%	

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $^{\dagger}$ Maine Youth Tobacco Survey, 2000.

# **Disparities Among Adult Population Groups, 2000\***





Maine is investing 125% of CDC's Best Practices lower estimated recommended funding and 55% of the upper estimated recommended funding. Maine spent \$377 per capita on smoking-attributable direct medical expenditures. In 1998, about 16% (\$169,000,000 or \$996.71 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

# **Smoking-Attributable Direct Medical Expenditures, 1998**

#### Ambulatory \$134,000,000 \$110,000,000 Hospital Nursing Home \$134,000,000 Prescription Drugs \$43,000,000 Other \$49,000,000 Annual Total \$470,000,000 Annual Per Capita \$377

# Smoking-Attributable **Productivity Costs, 1999**

Annual Total Annual Per Capita	\$406,000,000 \$324

### State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$47,268,164 \$77,235,000 Gross cigarette tax revenue collected in 2000 Cigarette tax per pack was \$0.74 in 2001

Cigarette sales were 82.9 packs per capita in 2000

FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)*	\$12,655,488	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$12,655,488	
Federal—CDC Office on Smoking and Health	\$1,001,691	6/01–5/02
Federal—SAMHSA	\$30,000	10/01-9/02
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	6/01–5/02	
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		\$14,018,672
CDC Best Practices Recommended Annual Total (Lower F	\$11,189,000	
CDC Best Practices Recommended Annual Total (Upper E	\$25,353,000	
FY02 Per Capita Investment in Tobacco Control	\$10.96	


<sup>\*</sup>An additional \$1,100,000 was appropriated to the Department of Health Services, Medical Care for payments to Medicaid health care providers for services that include smoking cessation and other health services.




SMOKING-ATTRIBUTABLE DEATHS, 1999		Smoking-Attributable Deaths, 1999 Disease-Specific		Youth Projected to Die from Smoking, 1999–2000	
Overall Men Women Death Rate	6,831 3,987 2,844 283.3/100,000	State Lung Cancer All States State CHD* All States	93.2/100,000 90.2/100,000 59.8/100,000 59.7/100,000	Overall Projected Death Rate Projected All	104,889 7,734/100,000
All States	295.5/100,000	State COPD† All States	61.2/100,000 59.7/100,000	States	8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

# Adult Cigarette Use, 2000



# **Environmental Tobacco Smoke, 1998–99**




#### **Youth Tobacco Use**

Grades 6–8			Grades 9–12		
Current Cigarette Smoking		Current Any Tobacco Use	Current Cigarette Current Any Smoking Tobacco Use		
National* MD <sup>†</sup> Boys <sup>†</sup> Girls <sup>†</sup>	11.0% 7.3% 7.3% 7.3%	15.1% 11.8% 12.5% 10.8%	28.0% 23.7% 23.3% 23.8%	34.5% 29.9% 32.1% 27.4%	

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $^{\dagger}$ Maryland Youth Tobacco Survey, 2000.

# **Disparities Among Adult Population Groups, 2000\***





Maryland is investing 107% of CDC's Best Practices lower estimated recommended funding and 41% of the upper estimated recommended funding. Maryland spent \$299 per capita on smoking-attributable direct medical expenditures. In 1998, about 15% (\$372,000,000 or \$663.31 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

# Smoking-Attributable Direct Medical Expenditures, 1998

# Ambulatory \$588,000,000 Hospital \$309,000,000 Nursing Home \$400,000,000 Prescription Drugs \$129,000,000 Other \$107,000,000 Annual Total \$1,533,000,000 Annual Per Capita \$299

# Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$1,555,000,000 \$301

### **State Revenue from Tobacco Sales and Settlement**

Tobacco settlement revenue received in 2001 \$140,202,664 Gross cigarette tax revenue collected in 2000 \$200,760,000

Cigarette tax per pack was \$0.66 in 2001

Cigarette sales were 57.7 packs per capita in 2000

Funding Source	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)*	\$30,671,765	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$ 30,671,765	
Federal—CDC Office on Smoking and Health	\$1,370,605	6/01–5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$257,036	6/01–5/02
Subtotal: Federal/National Sources	\$1,627,641	
FY02 Total Investment in Tobacco Control		\$32,299,406
CDC Best Practices Recommended Annual Total (Lower E	\$30,301,000	
CDC Best Practices Recommended Annual Total (Upper E	\$78,601,000	
FY02 Per Capita Investment in Tobacco Control	\$6.04	


<sup>\*</sup>A total of \$10,674,000 of the \$30,671,756 may not be expended until the Secretary of Budget and Management determines that sufficient tobacco settlement funds are available. An additional \$1,177,408 is budgeted by the Department of Health from general fund revenue for tobacco control.




Smoking-Attributable Deaths, 1999		Smoking-Attribut 1999 Disease-Sh	)	Youth Projected to Die from Smoking, 1999–2000	
Overall Men	9,008 5,064	State Lung Cancer All States	85.1/100,000 90.2/100,000	Overall Projected Death	117,753
Women Death Rate	3,944 261.2/100,000	State CHD* All States	47.2/100,000 59.7/100,000	Rate Projected All	7,850/100,000
All States	295.5/100,000	State COPD† All States	60.9/100,000 59.7/100,000	Štates	8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

#### **Adult Cigarette Use, 2000**



# **Environmental Tobacco Smoke, 1998-99**




#### **Youth Tobacco Use**

GRADES 6–8			GRADES 9–12		
Current Cigarette		Current Any	Current Cigarette	Current Any	
Smoking		Tobacco Use	Smoking	Tobacco Use	
National*	11.0%	15.1%	28.0%	34.5%	
MA <sup>†</sup>	n/a	n/a	30.3%	35.1%	
Boys <sup>†</sup>	n/a	n/a	29.9%	37.8%	
Girls <sup>†</sup>	n/a	n/a	30.7%	32.3%	

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000; †Massachusetts Youth Risk Behavior Survey, 1999.

# **Disparities Among Adult Population Groups, 2000\***





As of November 30, 2001, the state legislature had not completed its FY2002 budget bill; therefore, investment in tobacco control figures were not available for this report. Massachusetts spent \$450 per capita on smoking-attributable direct medical expenditures. In 1998, about 14% (\$817,000,000 or \$899.73 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

# Smoking-Attributable Direct Medical Expenditures, 1998

# Ambulatory \$861,000,000 Hospital \$570,000,000 Nursing Home \$934,000,000 Prescription Drugs \$194,000,000 Other \$207,000,000 Annual Total \$2,766,000,000 Annual Per Capita \$450

# Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$1,588,000,000 \$257

#### State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$240,246,288 Gross cigarette tax revenue collected in 2000 \$274,467,000

Cigarette tax per pack was \$0.76 in 2001

Cigarette sales were 58.3 packs per capita in 2000


Funding Source	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	not available	
State Appropriation—Excise Tax Revenue	not available	
State Appropriation—Other	not available	
Subtotal: State Appropriation	not available	
Federal—CDC Office on Smoking and Health	\$1,571,990	6/01–5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$449,370	6/01–5/02
Subtotal: Federal/National Sources	\$2,021,360	
FY02 Total Investment in Tobacco Control		not available
CDC Best Practices Recommended Annual Total (Lower	\$35,244,000	
CDC Best Practices Recommended Annual Total (Upper	\$92,758,000	
FY02 Per Capita Investment in Tobacco Control	not available	




SMOKING-ATTRIBUTABLE DEATHS, 1999		SMOKING-ATTRIBUT 1999 DISEASE-SF		YOUTH PROJECTED TO DIE FROM SMOKING, 1999–2000	
Overall Men Women Death Rate	14,741 8,905 5,836 299.0/100,000	State Lung Cancer All States State CHD* All States	88.6/100,000 90.2/100,000 71.3/100,000 59.7/100,000	Overall Projected Death Rate Projected All	255,356 9,837/100,000
All States	295.5/100,000	State COPD† All States	66.6/100,000 59.7/100,000	States	8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

# Adult Cigarette Use, 2000



#### **Environmental Tobacco Smoke, 1998-99**




#### **Youth Tobacco Use**

GRADES 6–8			Grades 9–12		
Current Cigarette Smoking		Current Any Tobacco Use	Current Cigarette Current Any Smoking Tobacco Use		
National* MI <sup>†</sup> Boys <sup>†</sup> Girls <sup>†</sup>	11.0% 9.3% 9.8% 8.7%	15.1% 14.2% 15.8% 12.4%	28.0% 27.6% 25.8% 27.5%	34.5% 34.1% 36.1% 32.0%	

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $^{\dagger}$ Michigan Youth Tobacco Survey, 2001.

# **Disparities Among Adult Population Groups, 2000\***





Michigan is investing 12% of CDC's Best Practices lower estimated recommended funding and 4% of the upper estimated recommended funding. Michigan spent \$270 per capita on smoking-attributable direct medical expenditures. In 1998, about 16% (\$881,000,000 or \$646.11 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

# Smoking-Attributable Direct Medical Expenditures, 1998

# Ambulatory \$897,000,000 Hospital \$734,000,000 Nursing Home \$575,000,000 Prescription Drugs \$286,000,000 Other \$163,000,000 Annual Total \$2,655,000,000 Annual Per Capita \$270

# Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$3,405,000,000 \$345

#### State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$258,862,265 Gross cigarette tax revenue collected in 2000 \$601,780,000

Cigarette tax per pack was \$0.75 in 2001

Cigarette sales were 83.7 packs per capita in 2000

Funding Source	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$0	
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other*	\$3,900,000	10/01–9/02
Subtotal: State Appropriation	\$3,900,000	
Federal—CDC Office on Smoking and Health	\$1,775,695	6/01–5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$318,530	10/00–9/01
Non-Government Source—RWJF/AMA	\$340,380	6/01–5/02
Subtotal: Federal/National Sources	\$2,434,605	
FY02 Total Investment in Tobacco Control		\$6,334,605
CDC Best Practices Recommended Annual Total (Lower 1	\$54,804,000	
CDC Best Practices Recommended Annual Total (Upper l	\$154,558,000	
FY02 Per Capita Investment in Tobacco Control		\$0.63


<sup>\*</sup>Funding amounts not verified by state budget office.




SMOKING-ATTRIBUTABLE DEATHS, 1999	SMOKING-ATTRIBUT 1999 DISEASE-SI	<b>)</b>	YOUTH PROJECTED TO DIE FROM SMOKING, 1999–2000	
Overall         5,618           Men         3,410           Women         2,208           Death Rate         231.1/100,000           All States         295.5/100,000	State Lung Cancer All States State CHD* All States State COPD† All States	73.8/100,000 90.2/100,000 37.9/100,000 59.7/100,000 58.9/100,000 59.7/100,000	Overall Projected Death Rate Projected All States	112,085 8,710/100,000 8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

# **Adult Cigarette Use, 2000**



# **Environmental Tobacco Smoke, 1998-99**




#### **Youth Tobacco Use**

Grades 6–8			GRADES 9–12	
Current Cigarette Smoking		Current Any Tobacco Use	Current Cigarette Current A Smoking Tobacco	
National* MN <sup>†</sup> Boys <sup>†</sup> Girls <sup>†</sup>	11.0% 9.1% 8.7% 9.5%	15.1% 12.6% 12.9% 12.3%	28.0% 32.4% 32.0% 32.6%	34.5% 38.7% 42.7% 34.1%

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $^{\dagger}$ Minnesota Youth Tobacco Survey, 2000.

#### Disparities Among Adult Population Groups, 2000\*





Minnesota is investing 105% of CDC's Best Practices lower estimated recommended funding and 41% of the upper estimated recommended funding. Minnesota spent \$341 per capita on smoking-attributable direct medical expenditures. In 1998, about 13% (\$363,000,000 or \$675.57 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

# **Smoking-Attributable Direct Medical Expenditures, 1998**

Ambulatory Hospital Nursing Home Prescription Drugs Other Annual Total	\$679,000,000 \$296,000,000 \$434,000,000 \$105,000,000 \$97,000,000 \$1,611,000,000
Annual Total Annual Per Capita	\$1,611,000,000 \$341
•	

### Smoking-Attributable **Productivity Costs, 1999**

Annual Total	\$1,027,000,000
Annual Per Capita	\$215

# State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 Gross cigarette tax revenue collected in 2000 \$336,997,000

\$177,262,000

Cigarette tax per pack was \$0.48 in 2001


Cigarette sales were 76.0 packs per capita in 2000

FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$17,610,935	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Non-Governmental Funding—MN Partnership for Action Against Tobacco	\$10,900,000	7/01–6/02
Subtotal: State Sources	\$28,510,935	
Federal—CDC Office on Smoking and Health	\$1,376,165	6/01–5/02
Federal—SAMHSA	\$71,710	10/01-9/02
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$232,748	03/01–2/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control	\$30,191,558	
CDC Best Practices Recommended Annual Total (Lower E	\$28,624,000	
CDC Best Practices Recommended Annual Total (Upper E	\$74,013,000	
FY02 Per Capita Investment in Tobacco Control	<b>\$6.07</b>	


SMOKING-ATTRIBUTABLE DEATHS, 1999		Smoking-Attributable Deaths, 1999 Disease-Specific		Youth Projected to Die from Smoking, 1999–2000	
Overall Men	4,961 3,306	State Lung Cancer All States	115.0/100,000 90.2/100,000	Overall Projected Death	58,515
Women Death Rate	1,655 367.8/100,000	State CHD* All States	80.2/100,000 59.7/100,000	Rate Projected All	7,549/100,000
All States	295.5/100,000	State COPD† All States	71.5/100,000 59.7/100,000	Štates	8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

# Adult Cigarette Use, 2000



# Environmental Tobacco Smoke, 1998–99




#### **Youth Tobacco Use**

Grades 6–8			GRADES 9–12	
Current Cigarette Smoking		Current Any Tobacco Use	Current Cigarette Current Smoking Tobacco	
National* MS <sup>†</sup> Boys <sup>†</sup> Girls <sup>†</sup>	11.0% 17.8% 19.2% 16.3%	15.1% 25.3% 29.4% 20.8%	28.0% 30.5% 33.3% 27.6%	34.5% 40.6% 47.8% 33.3%

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $^{\dagger}$ Mississippi Youth Tobacco Survey, 2000.

# **Disparities Among Adult Population Groups, 2000\***





Mississippi is investing 120% of CDC's Best Practices lower estimated recommended funding and 48% of the upper estimated recommended funding. Mississippi spent \$204 per capita on smoking-attributable direct medical expenditures. In 1998, about 14% (\$206,000,000 or \$424.29 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

# **Smoking-Attributable Direct Medical Expenditures, 1998**

Ambulatory Hospital Nursing Home Prescription Drugs Other Annual Total	\$184,000,000 \$154,000,000 \$130,000,000 \$55,000,000 \$38,000,000 \$561,000,000
Annual Per Capita	\$204

# Smoking-Attributable **Productivity Costs, 1999**

Annual Total Annual Per Capita	\$1,298,000,000 \$469

#### **State Revenue from Tobacco Sales and Settlement**

Tobacco settlement revenue received in 2001

\$211,149,286

Gross cigarette tax revenue collected in 2000

\$49,247,000

Cigarette tax per pack was \$0.18 in 2001


Cigarette sales were 97.2 packs per capita in 2000

FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$0	
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Non-Governmental Source—Partnership for a Healthier	\$21,782,613	
Mississippi		
Subtotal: State Sources	\$21,782,613	
Federal—CDC Office on Smoking and Health	\$380,796	6/01–5/02
Federal—SAMHSA	\$131,834	10/01–9/02
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA \$299,196		6/01–5/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control	\$22,594,439	
CDC Best Practices Recommended Annual Total (Lower E	\$18,788,000	
CDC Best Practices Recommended Annual Total (Upper E	\$46,804,000	
FY02 Per Capita Investment in Tobacco Control	\$7.87	


Smoking-Attributable Deaths, 1999		SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
Overall Men Women Death Rate	10,271 6,238 4,033 345.3/100,000	State Lung Cancer All States State CHD* All States	101.6/100,000 90.2/100,000 79.7/100,000 59.7/100,000	Overall Projected Death Rate Projected All	139,484 9,770/100,000
All States	295.5/100,000	State COPD† All States	78.9/100,000 59.7/100,000	Štates	8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

# Adult Cigarette Use, 2000



#### **Environmental Tobacco Smoke, 1998-99**




#### **Youth Tobacco Use**

Grades 6–8			Grades 9–12	
Current Cigarette Smoking		Current Any Tobacco Use	Current Cigarette Current A Smoking Tobacco	
National* MO Boys Girls	11.0% 14.9%† 13.7%† 16.3%†	15.1% 19.5%† n/a n/a	28.0% 32.8%†† 35.6%†† 30.1%††	34.5% 39.0%†† 44.0%†† 33.9%††

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000; †Missouri Youth Tobacco Survey, 1999; ††Missouri Youth Risk Behavior Survey, 1999.

# **Disparities Among Adult Population Groups, 2000\***





Missouri is investing 72% of CDC's *Best Practices* lower estimated recommended funding and 26% of the upper estimated recommended funding. Missouri spent \$307 per capita on smoking-attributable direct medical expenditures. In 1998, about 14% (\$415,000,000 or \$565.59 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

# Smoking-Attributable Direct Medical Expenditures, 1998

# Ambulatory \$507,000,000 Hospital \$452,000,000 Nursing Home \$466,000,000 Prescription Drugs \$133,000,000 Other \$110,000,000 Annual Total \$1,668,000,000 Annual Per Capita \$307

# Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$2,173,000,000 \$397

# State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$142,110,326 Gross cigarette tax revenue collected in 2000 \$107,169,000

Cigarette tax per pack was \$0.17 in 2001

Cigarette sales were 113.8 packs per capita in 2000

FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)*	\$22,176,700	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$22,176,700	
Federal—CDC Office on Smoking and Health	\$1,166,052	6/01–5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$200,000	6/01–5/02
Subtotal: Federal/National Sources	\$1,366,052	
FY02 Total Investment in Tobacco Control		\$23,542,752
CDC Best Practices Recommended Annual Total (Lower F	Estimate)	\$32,767,000
CDC Best Practices Recommended Annual Total (Upper F	Estimate)	\$91,359,000
FY02 Per Capita Investment in Tobacco Control		\$4.17


<sup>\*</sup>Due to a budget shortfall, \$3,500,000 of this money is withheld.




SMOKING-ATTRIBUTABL 1999	E DEATHS,	Smoking-Attribut 1999 Disease-Sp		Youth Pr to Die from 1999–	SMOKING,
	1,434 858 576 7/100,000 5/100,000	State Lung Cancer All States State CHD* All States State COPD† All States	86.3/100,000 90.2/100,000 44.5/100,000 59.7/100,000 90.3/100,000 59.7/100,000	Overall Projected Death Rate Projected All States	17,143 7,451/100,000 8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

# Adult Cigarette Use, 2000



# **Environmental Tobacco Smoke, 1998–99**




#### **Youth Tobacco Use**

Grades 6–8		GRADES 9–12		
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
MT <sup>†</sup>	n/a	n/a	35.0%	45.0%
Boys <sup>†</sup>	n/a	n/a	35.4%	52.3%
Girls <sup>†</sup>	n/a	n/a	34.6%	37.6%

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $^{\dagger}$ Montana Youth Risk Behavior Survey, 1999.

# **Disparities Among Adult Population Groups, 2000\***





Montana is investing 20% of CDC's *Best Practices* lower estimated recommended funding and 10% of the upper estimated recommended funding. Montana spent \$246 per capita on smoking-attributable direct medical expenditures. In 1998, about 13% (\$52,000,000 or \$514.32 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

# Smoking-Attributable Direct Medical Expenditures, 1998

# Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$247,000,000 \$280

#### State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$26,248,117 Gross cigarette tax revenue collected in 2000 \$12,442,000

Cigarette tax per pack was \$0.18 in 2001


Cigarette sales were 75.5 packs per capita in 2000

FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$0	
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$502,165	7/01–6/02
Subtotal: State Appropriation	\$502,165	
Federal—CDC Office on Smoking and Health	\$875,000	6/01–5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$178,282	10/00–9/01
Non-Government Source—RWJF/AMA	\$349,174	6/01–5/02
Subtotal: Federal/National Sources	\$1,402,456	
FY02 Total Investment in Tobacco Control		\$1,904,621
CDC Best Practices Recommended Annual Total (Lower E	stimate)	\$9,355,000
CDC Best Practices Recommended Annual Total (Upper E	stimate)	\$19,679,000
FY02 Per Capita Investment in Tobacco Control		\$2.09


	IBUTABLE DEATHS, 1999	Smoking-Attribut 1999 Disease-Sh	)	TO DIE FRO	Projected om Smoking, 9–2000
Overall Men Women	2,449 1,552 897	State Lung Cancer All States State CHD*	81.2/100,000 90.2/100,000 43.5/100,000	Overall Projected Death Rate	45,712 h 10,153/100,000
Death Rate All States	264.2/100,000 295.5/100,000	All States State COPD† All States	59.7/100,000 72.7/100,000 59.7/100,000	Projected All States	8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

# Adult Cigarette Use, 2000



# **Environmental Tobacco Smoke, 1998–99**




#### **Youth Tobacco Use**

GRADES 6-8		Grades 9–12		
Current C		Current Any	Current Cigarette	Current Any
Smol		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
NE	10.0%†	13.8% <sup>†</sup>	29.0%††	35.8%††
Boys	10.2%†	15.6% <sup>†</sup>	27.1%††	38.9%††
Girls	9.8%†	11.8% <sup>†</sup>	30.8%††	32.8%††

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $\dagger$ Nebraska Youth Tobacco Survey, 1999;  $\dagger$  $\dagger$ Nebraska Youth Tobacco Survey, 2000.

#### **Disparities Among Adult Population Groups, 2000\***





Nebraska is investing 65% of CDC's *Best Practices* lower estimated recommended funding and 28% of the upper estimated recommended funding. Nebraska spent \$252 per capita on smoking-attributable direct medical expenditures. In 1998, about 12% (\$105,000,000 or \$499.41 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

# Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory Hospital Nursing Home Prescription Drugs Other Annual Total	\$116,000,000 \$104,000,000 \$139,000,000 \$39,000,000 \$21,000,000 \$419,000,000
Annual Per Capita	\$252

# Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$439,000,000 \$264

#### State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$36,903,219 Gross cigarette tax revenue collected in 2000 \$44,994,000

Cigarette tax per pack was \$0.34 in 2001

Cigarette sales were 77.6 packs per capita in 2000


Funding Source	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$7,000,000	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$7,000,000	
Federal—CDC Office on Smoking and Health	\$1,347,138	6/01–5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$296,098	6/01–5/02
Subtotal: Federal/National Sources	\$1,643,236	
FY02 Total Investment in Tobacco Control		\$8,643,236
CDC Best Practices Recommended Annual Total (Lower E	stimate)	\$13,308,000
CDC Best Practices Recommended Annual Total (Upper E	stimate)	\$31,041,000
FY02 Per Capita Investment in Tobacco Control		\$5.01




Overall 3,359 State Lung Cancer 113.0/100,000 Overall	
Men         1,910         All States         90.2/100,000         Projected Death           Women         1,449         State CHD*         82.7/100,000         Rate         11,08           Death Rate         414.3/100,000         All States         59.7/100,000         Projected Death	56,725 4/100,000 0/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

# Adult Cigarette Use, 2000



# **Environmental Tobacco Smoke, 1998-99**




#### **Youth Tobacco Use**

Grades 6–8			GRADES 9–12	
Current Cigarette		Current Any	Current Cigarette	Current Any
Smoking		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
NV <sup>†</sup>	n/a	n/a	32.6%	39.9%
Boys <sup>†</sup>	n/a	n/a	32.4%	44.6%
Girls <sup>†</sup>	n/a	n/a	32.5%	34.5%

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000; <sup>†</sup>Nevada Youth Risk Behavior Survey, 1999.

#### Disparities Among Adult Population Groups, 2000\*





Nevada is investing 38% of CDC's *Best Practices* lower estimated recommended funding and 16% of the upper estimated recommended funding. Nevada spent \$253 per capita on smoking-attributable direct medical expenditures. In 1998, about 19% (\$96,000,000 or \$748.56 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

# Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory Hospital Nursing Home Prescription Drugs Other Annual Total	\$199,000,000 \$113,000,000 \$43,000,000 \$41,000,000 \$45,000,000 \$441,000,000
Annual Per Capita	\$253

# Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$762,000,000 \$421

# State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$37,830,589 Gross cigarette tax revenue collected in 2000 \$61,017,000

Cigarette tax per pack was \$0.35 in 2001


Cigarette sales were 93.2 packs per capita in 2000

Funding Source	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$4,093,565	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other		
Subtotal: State Appropriation	\$4,093,565	
Federal—CDC Office on Smoking and Health	\$748,437	6/01–5/02
Federal—SAMHSA	10/01–9/02	
Non-Government Source—American Legacy Foundation		
Non-Government Source—RWJF/AMA	6/01–5/02	
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		\$5,116,564
CDC Best Practices Recommended Annual Total (Lower E	\$13,477,000	
CDC Best Practices Recommended Annual Total (Upper E	\$32,993,000	
FY02 Per Capita Investment in Tobacco Control	\$2.46	


SMOKING-ATTRIBUTABLE DEATHS, 1999	SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		YOUTH PROJECTED TO DIE FROM SMOKING, 1999–2000	
Overall       1,688         Men       989         Women       699         Death Rate       291.9/100,000         All States       295.5/100,000	State Lung Cancer All States State CHD* All States State COPD† All States	92.5/100,000 90.2/100,000 58.4/100,000 59.7/100,000 78.7/100,000 59.7/100,000	Overall Projected Deatl Rate Projected All States	34,700 1 11,209/100,000 8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

# Adult Cigarette Use, 2000



# **Environmental Tobacco Smoke, 1998–99**




#### **Youth Tobacco Use**

GRADES 6–8		GRADES 9–12		
Current Cigarette		Current Any	Current Cigarette	Current Any
Smoking		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
NH	12.0% <sup>†</sup>	15.2% <sup>†</sup>	36.0% <sup>††</sup>	n/a
Boys	11.1% <sup>†</sup>	16.0% <sup>†</sup>	32.0% <sup>††</sup>	n/a
Girls	12.8% <sup>†</sup>	14.2% <sup>†</sup>	39.9% <sup>††</sup>	n/a

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000; <sup>†</sup>New Hampshire Youth Tobacco Survey, 2000; <sup>††</sup>New Hampshire Youth Risk Behavior Survey, 1995.

# **Disparities Among Adult Population Groups, 2000\***





New Hampshire is investing 40% of CDC's Best Practices lower estimated recommended funding and 18% of the upper estimated recommended funding. New Hampshire spent \$371 per capita on smoking-attributable direct medical expenditures. In 1998, about 13% (\$90,000,000 or \$958.66 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

# Smoking-Attributable Direct Medical Expenditures, 1998

# Ambulatory \$153,000,000 Hospital \$90,000,000 Nursing Home \$119,000,000 Prescription Drugs \$37,000,000 Other \$41,000,000 Annual Total \$440,000,000 Annual Per Capita \$371

# Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$339,000,000 \$282

# State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$43,303,862 Gross cigarette tax revenue collected in 2000 \$93,588,000

Cigarette tax per pack was \$0.52 in 2001


Cigarette sales were 147.3 packs per capita in 2000

FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$3,000,000	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$3,000,000	
Federal—CDC Office on Smoking and Health	\$991,588	6/01–5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$414,957	6/01–5/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		\$4,406,545
CDC Best Practices Recommended Annual Total (Lower E	\$10,888,000	
CDC Best Practices Recommended Annual Total (Upper E	\$24,766,000	
FY02 Per Capita Investment in Tobacco Control	\$2.33	


SMOKING-ATTRIBUTABLE DEATHS, 1999	SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		YOUTH PROJECTED TO DIE FROM SMOKING, 1999–2000	
Overall       10,756         Men       6,332         Women       4,424         Death Rate       243.7/100,000         All States       295.5/100,000	All States State CHD* All States	80.7/100,000 90.2/100,000 53.5/100,000 59.7/100,000 51.1/100,000 59.7/100,000	Overall Projected Death Rate Projected All States	168,828 8,087/100,000 8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

# Adult Cigarette Use, 2000



#### **Environmental Tobacco Smoke, 1998-99**




#### **Youth Tobacco Use**

Grades 6–8		GRADES 9–12		
Current Cigarette		Current Any	Current Cigarette	Current Any
Smoking		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
NJ <sup>†</sup>	10.5%	18.9%	27.6%	38.9%
Boys <sup>†</sup>	9.9%	20.4%	26.4%	42.3%
Girls <sup>†</sup>	11.0%	17.3%	28.4%	35.0%

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $^{\dagger}$ New Jersey Youth Tobacco Survey, 1999.

# Disparities Among Adult Population Groups, 2000\*





New Jersey is investing 72% of CDC's Best Practices lower estimated recommended funding and 27% of the upper estimated recommended funding. New Jersey spent \$306 per capita on smoking-attributable direct medical expenditures. In 1998, about 16% (\$755,000,000 or \$928.76 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

# Smoking-Attributable Direct Medical Expenditures, 1998

# Ambulatory \$854,000,000 Hospital \$497,000,000 Nursing Home \$712,000,000 Prescription Drugs \$255,000,000 Other \$163,000,000 Annual Total \$2,481,000,000 Annual Per Capita \$306

# Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$2,226,000,000 \$273

# State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$239,846,695 Gross cigarette tax revenue collected in 2000 \$393,179,000

Cigarette tax per pack was \$0.80 in 2001


Cigarette sales were 60.1 packs per capita in 2000

Funding Source	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$30,000,000	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$30,000,000	
Federal—CDC Office on Smoking and Health	\$1,257,351	6/01–5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$757,118	10/00–9/01
Non-Government Source—RWJF/AMA	\$447,197	3/01–2/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		\$32,461,666
CDC Best Practices Recommended Annual Total (Lower E	\$45,073,000	
CDC Best Practices Recommended Annual Total (Upper E	\$121,328,000	
FY02 Per Capita Investment in Tobacco Control	\$3.83	


SMOKING-ATTRIBUTABLE DEATHS, 1999		SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
	2,109 1,302 807 4.7/100,000 5.5/100,000	State Lung Cancer All States State CHD* All States State COPD† All States	62.8/100,000 90.2/100,000 55.2/100,000 59.7/100,000 82.3/100,000 59.7/100,000	Overall Projected Death Rate Projected All States	44,564 8,763/100,000 8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

#### **Adult Cigarette Use, 2000**



# **Environmental Tobacco Smoke, 1998-99**




#### **Youth Tobacco Use**

Grades 6–8		Grades 9–12		
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
NM <sup>†</sup>	n/a	n/a	30.1%	n/a
Boys <sup>†</sup>	n/a	n/a	n/a	n/a
Girls <sup>†</sup>	n/a	n/a	n/a	n/a

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $^{\dagger}$ New Mexico Youth Risk Behavior Survey, 1991.

# **Disparities Among Adult Population Groups, 2000\***





New Mexico is investing 51% of CDC's *Best Practices* lower estimated recommended funding and 22% of the upper estimated recommended funding. New Mexico spent \$208 per capita on smoking-attributable direct medical expenditures. In 1998, about 15% (\$144,000,000 or \$436.51 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

# Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory Hospital Nursing Home Prescription Drugs Other Annual Total	\$135,000,000 \$103,000,000 \$58,000,000 \$29,000,000 \$35,000,000
Annual Per Capita	\$208

# Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$397,000,000 \$228

# State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$36,989,850 Gross cigarette tax revenue collected in 2000 \$21,007,000

Cigarette tax per pack was \$0.21 in 2001

Cigarette sales were 53.8 packs per capita in 2000


Funding Source	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$5,000,000	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other		
Subtotal: State Appropriation	\$5,000,000	
Federal—CDC Office on Smoking and Health	\$1,181,300	6/01–5/02
Federal—SAMHSA		
Non-Government Source—American Legacy Foundation	\$460,229	10/00–9/01
Non-Government Source—RWJF/AMA	\$293,480	6/01–5/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control	\$6,935,009	
CDC Best Practices Recommended Annual Total (Lower E	\$13,711,000	
CDC Best Practices Recommended Annual Total (Upper E	\$31,947,000	
FY02 Per Capita Investment in Tobacco Control	\$3.75	




SMOKING-ATTRIBUTABLE DEATHS, 1999		SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
Overall Men Women Death Rate	24,572 13,439 11,133 252.1/100,000	State Lung Cancer All States State CHD* All States	76.7/100,000 90.2/100,000 66.7/100,000 59.7/100,000	Overall Projected Death Rate Projected All	438,769 9,355/100,000
All States	295.5/100,000	State COPD† All States	51.5/100,000 59.7/100,000	States	8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

# Adult Cigarette Use, 2000



#### **Environmental Tobacco Smoke, 1998–99**




#### **Youth Tobacco Use**

GRADES 6–8		GRADES 9–12		
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
NY <sup>†</sup>	9.3%	11.8%	26.8%	32.8%
Boys <sup>†</sup>	9.8%	13.5%	24.8%	34.2%
Girls <sup>†</sup>	8.8%	9.9%	29.2%	30.9%

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $^{\dagger}$ New York Youth Tobacco Survey, 2000.

# Disparities Among Adult Population Groups, 2000\*





New York is investing 49% of CDC's Best Practices lower estimated recommended funding and 17% of the upper estimated recommended funding. New York spent \$351 per capita on smoking-attributable direct medical expenditures. In 1998, about 16% (\$4,271,000,000 or \$1,389.77 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

# Smoking-Attributable Direct Medical Expenditures, 1998

# Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$5,303,000,000 \$291

### State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$754,339,700 Gross cigarette tax revenue collected in 2000 \$738,115,000

Cigarette tax per pack was \$1.11 in 2001 and will increase to \$1.50 effective April 2002

Cigarette sales were 57.8 packs per capita in 2000

FUNDING SOURCE	FUNDING CYCLE	
State Appropriation—Settlement (Tobacco Only)*	\$42,500,000	4/01–3/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation		
Federal—CDC Office on Smoking and Health	\$1,999,998	6/01–5/02
Federal—SAMHSA		
Non-Government Source—American Legacy Foundation	10/00–9/01	
Non-Government Source—RWJF/AMA	6/01–5/02	
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control	\$46,479,998	
CDC Best Practices Recommended Annual Total (Lower	\$95,830,000	
CDC Best Practices Recommended Annual Total (Upper l	\$269,296,000	
FY02 Per Capita Investment in Tobacco Control	\$2.44	


<sup>\*</sup>An additional \$2,000,000 is appropriated to the Department of Health for administrative costs for cancer and tobacco control programs. Also, an additional \$2,000,000 was appropriated for services and expenses related to tobacco control and cancer programs. In New York State, approximately one-half of all Tobacco Settlement funds are shared with counties. These figures represent the New York State share of Tobacco Settlement appropriations and do not include settlement funds appropriated by counties for tobacco prevention and control.




Overall         11,511         State Lung Cancer         96.6/100,000         Overall         21           Men         7,388         All States         90.2/100,000         Projected Death	
Women	

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

#### **Adult Cigarette Use, 2000**



# **Environmental Tobacco Smoke, 1998-99**




#### **Youth Tobacco Use**

GRADES 6–8		GRADES 9–12		
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
NC <sup>†</sup>	15.0%	18.4%	31.6%	38.3%
Boys <sup>†</sup>	16.0%	21.0%	33.4%	44.0%
Girls <sup>†</sup>	14.0%	15.7%	29.7%	32.4%

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $^{\dagger}$ North Carolina Youth Tobacco Survey, 1999.

# **Disparities Among Adult Population Groups, 2000\***





North Carolina is investing 9% of CDC's *Best Practices* lower estimated recommended funding and 3% of the upper estimated recommended funding. North Carolina spent \$255 per capita on smoking-attributable direct medical expenditures. In 1998, about 13% (\$600,000,000 or \$513.30 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

# Smoking-Attributable Direct Medical Expenditures, 1998

# Ambulatory \$646,000,000 Hospital \$478,000,000 Nursing Home \$491,000,000 Prescription Drugs \$161,000,000 Other \$147,000,000 Annual Total \$1,923,000,000 Annual Per Capita \$255

# Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$2,829,000,000 \$370

# State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001

\$142,728,818 \$42,378,000

Gross cigarette tax revenue collected in 2000

Cigarette tax per pack was \$0.05 in 2001


Cigarette sales were 109.0 packs per capita in 2000

FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$0	
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$650,000	7/01–6/02
Subtotal: State Appropriation	\$650,000	
Federal—CDC Office on Smoking and Health	\$1,848,922	6/01–5/02
Federal—SAMHSA	\$208,961	10/01–9/02
Non-Government Source—American Legacy Foundation	\$1,141,057	10/00–9/01
Non-Government Source—RWJF/AMA	\$0	
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		\$3,848,940
CDC Best Practices Recommended Annual Total (Lower E	\$42,591,000	
CDC Best Practices Recommended Annual Total (Upper E	\$118,626,000	
FY02 Per Capita Investment in Tobacco Control	\$0.47	


Smoking-Attributable Deaths, 1999	SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		YOUTH PROJECTED TO DIE FROM SMOKING, 1999–2000	
Overall       855         Men       593         Women       262         Death Rate       227.7/100,000         All States       295.5/100,000	State Lung Cancer All States State CHD* All States State COPD† All States	64.1/100,000 90.2/100,000 53.1/100,000 59.7/100,000 50.9/100,000 59.7/100,000	Overall Projected Death Rate Projected All States	15,638 9,722/100,000 8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

# **Adult Cigarette Use, 2000**



# **Environmental Tobacco Smoke, 1998–99**




#### **Youth Tobacco Use**

GRADES 6–8		GRADES 9–12		
Current C Smok			Current Any Tobacco Use	
National* ND† Boys† Girls†	11.0% n/a n/a n/a	15.1% n/a n/a n/a	28.0% 40.6% 40.2% 41.0%	34.5% n/a n/a n/a

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $^{\dagger}$ North Dakota Youth Risk Behavior Survey, 1999.

# Disparities Among Adult Population Groups, 2000\*





North Dakota is investing 41% of CDC's Best Practices lower estimated recommended funding and 20% of the upper estimated recommended funding. North Dakota spent \$303 per capita on smoking-attributable direct medical expenditures. In 1998, about 11% (\$37,000,000 or \$604.68 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

# Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory Hospital Nursing Home Prescription Drugs Other Annual Total	\$54,000,000 \$56,000,000 \$61,000,000 \$13,000,000 \$9,000,000 \$193,000,000
Annual Per Capita	\$303

# Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$158,000,000 \$249

# State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$22,701,724 Gross cigarette tax revenue collected in 2000 \$21,132,000

Cigarette tax per pack was \$0.44 in 2001

Cigarette sales were 72.5 packs per capita in 2000


Funding Source	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$2,055,000	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$2,055,000	
Federal—CDC Office on Smoking and Health	\$1,099,997	6/01–5/02
Federal—SAMHSA	\$30,537	10/01-9/02
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$200,000	6/01–5/02
Subtotal: Federal/National Sources	\$1,330,534	
FY02 Total Investment in Tobacco Control	\$3,385,534	
CDC Best Practices Recommended Annual Total (Lower	\$8,161,000	
CDC Best Practices Recommended Annual Total (Upper	\$16,547,000	
FY02 Per Capita Investment in Tobacco Control	\$5.27	




SMOKING-ATTRIBUTABLE DEATHS, 1999		SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
Overall Men Women Death Rate	18,953 11,077 7,892 317.6/100,000	State Lung Cancer All States State CHD* All States	97.6/100,000 90.2/100,000 70.4/100,000 59.7/100,000	Overall Projected Deatl Rate Projected All	10,896/100,000
All States	295.5/100,000	State COPD† All States	74.2/100,000 59.7/100,000	Štates	8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

# Adult Cigarette Use, 2000



# **Environmental Tobacco Smoke, 1998–99**




#### **Youth Tobacco Use**

GRADES 6–8		GRADES 9–12		
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
OH <sup>†</sup>	13.7%	18.7%	33.4%	41.1%
Boys <sup>†</sup>	14.7%	22.0%	33.0%	45.6%
Girls <sup>†</sup>	12.7%	15.2%	33.7%	35.9%

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $\dagger$ Ohio Youth Tobacco Survey, 2000.

# **Disparities Among Adult Population Groups, 2000\***





Ohio is investing 197% of CDC's Best Practices lower estimated recommended funding and 70% the upper estimated recommended funding. Ohio spent \$304 per capita on smoking-attributable direct medical expenditures. In 1998, about 17% (\$1,113,000,000 or \$862.47 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

# **Smoking-Attributable Direct Medical Expenditures, 1998**

#### **Ambulatory** \$1,030,000,000 \$808,000,000 Hospital Nursing Home \$1,113,000,000 Prescription Drugs \$274,000,000 Other \$191,000,000 Annual Total \$3,416,000,000 Annual Per Capita \$304

# Smoking-Attributable **Productivity Costs, 1999**

Annual Total Annual Per Capita	\$4,146,000,000 \$368

#### State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$312,446,299 Gross cigarette tax revenue collected in 2000 \$271,260,000

Cigarette tax per pack was \$0.24 in 2001

Cigarette sales were 99.9 packs per capita in 2000


Funding Source	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)*	\$119,578,148	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$119,578,148	
Federal—CDC Office on Smoking and Health	\$1,525,232	6/01–5/02
Federal—SAMHSA	\$82,950	10/01–9/02
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$444,658	3/01–2/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control	\$121,630,988	
CDC Best Practices Recommended Annual Total (Lower E	\$61,735,000	
CDC Best Practices Recommended Annual Total (Upper E	\$173,676,000	
FY02 Per Capita Investment in Tobacco Control	\$10.67	

<sup>\*</sup>The amount reflected above is disbursed to an endowment fund where the funds will be used by the Foundation for tobacco use reduction program. The Foundation may, but is not required to, treat these funds as an endowment. An additional \$636,000 will be expended by the Department of Public Safety to enforce underage-use laws in establishments with liquor stores. Also, an additional \$579,323 will go to the Department of Health to provide emergency medications and oxygen to lowincome seniors adversely affected by tobacco.


Smoking-Attributable Deaths, 1999		Smoking-Attributable Deaths, 1999 Disease-Specific		Youth Projected to Die from Smoking, 1999–2000	
Overall Men Women Death Rate	5,781 3,589 2,192 320.1/100,000	State Lung Cancer All States State CHD* All States	98.4/100,000 90.2/100,000 75.3/100,000 59.7/100,000	Overall Projected Death Rate Projected All	77,943 8,734/100,000
All States	295.5/100,000	State COPD† All States	71.3/100,000 59.7/100,000	States	8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

# Adult Cigarette Use, 2000



#### **Environmental Tobacco Smoke, 1998–99**




#### **Youth Tobacco Use**

Grades 6–8		GRADES 9–12		
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
OK†	16.9%	21.0%	33.0%	42.0%
Boys†	17.9%	23.8%	31.8%	47.3%
Girls†	15.8%	17.8%	34.0%	36.4%

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $\dagger$ Oklahoma Youth Tobacco Survey, 1999.

# **Disparities Among Adult Population Groups, 2000\***





Oklahoma is investing 17% of CDC's *Best Practices* lower estimated recommended funding and 7% of the upper estimated recommended funding. Oklahoma spent \$272 per capita on smoking-attributable direct medical expenditures. In 1998, about 13% (\$170,000,000 or \$498.01 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

# Smoking-Attributable Direct Medical Expenditures, 1998

# Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$1,334,000,000 \$397

# State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001

\$64,265,818

Gross cigarette tax revenue collected in 2000

\$63,926,000

Cigarette tax per pack was \$0.23 in 2001

Cigarette sales were 108.9 packs per capita in 2000


Funding Source	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$1,750,000	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$1,750,000	
Federal—CDC Office on Smoking and Health	\$1,299,907	6/01–5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$750,000	10/00-9/01
Non-Government Source—RWJF/AMA	\$0	
Subtotal: Federal/National Sources	\$2,049,907	
FY02 Total Investment in Tobacco Control	\$3,799,907	
CDC Best Practices Recommended Annual Total (Lower E	\$21,825,000	
CDC Best Practices Recommended Annual Total (Upper E	\$56,310,000	
FY02 Per Capita Investment in Tobacco Control	\$1.09	




SMOKING-ATTRIBUTABLE DEATHS, 1999		Smoking-Attributable Deaths, 1999 Disease-Specific		Youth Projected to Die from Smoking, 1999–2000	
Overall Men Women Death Rate	4,986 2,991 1,995 278.6/100,000	State Lung Cancer All States State CHD* All States	85.7/100,000 90.2/100,000 47.2/100,000 59.7/100,000	Overall Projected Death Rate Projected All	73,132 8,639/100,000
All States	295.5/100,000	State COPD† All States	74.3/100,000 59.7/100,000	States	8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

# Adult Cigarette Use, 2000



# **Environmental Tobacco Smoke, 1998–99**




#### **Youth Tobacco Use**

GRADES 6–8		Grades 9–12		
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
OR	n/a	n/a	n/a	n/a
Boys	n/a	n/a	n/a	n/a
Girls	n/a	n/a	n/a	n/a

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Source: \*National Youth Tobacco Survey, 2000.

# **Disparities Among Adult Population Groups, 2000\***





Oregon is investing 62% of CDC's *Best Practices* lower estimated recommended funding and 25% of the upper estimated recommended funding. Oregon spent \$265 per capita on smoking-attributable direct medical expenditures. In 1998, about 13% (\$224,000,000 or \$438.97 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

# Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory Hospital Nursing Home Prescription Drugs Other Annual Total	\$328,000,000 \$182,000,000 \$208,000,000 \$74,000,000 \$79,000,000 \$871,000,000
Annual Per Capita	\$265

# Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$908,000,000 \$274

# State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$68,264,932 Gross cigarette tax revenue collected in 2000 \$166,359,000

Cigarette tax per pack was \$0.68 in 2001

Cigarette sales were 72.0 packs per capita in 2000


FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$2,500,000	7/01–6/02
State Appropriation—Excise Tax Revenue	\$9,086,500	7/01–6/02
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$11,586,500	
Federal—CDC Office on Smoking and Health	\$1,112,463	6/01–5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$353,453	6/01–5/02
Subtotal: Federal/National Sources	\$1,465,916	
FY02 Total Investment in Tobacco Control	\$13,052,416	
CDC Best Practices Recommended Annual Total (Lower F	\$21,131,000	
CDC Best Practices Recommended Annual Total (Upper F	\$52,840,000	
FY02 Per Capita Investment in Tobacco Control	\$3.75	




SMOKING-ATTRIBUTABLE DEATHS, 1999		Smoking-Attributable Deaths, 1999 Disease-Specific		Youth Projected to Die from Smoking, 1999–2000	
Overall Men Women	19,798 12,045 7,753	State Lung Cancer All States State CHD*	85.3/100,000 90.2/100,000 59.1/100.000	Overall Projected Deatl Rate	311,074 1 10,645/100,000
Death Rate All States	270.2/100,000 295.5/100,000	All States State COPD† All States	59.7/100,000 59.7/100,000 59.7/100,000	Projected All States	8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

# Adult Cigarette Use, 2000



# **Environmental Tobacco Smoke, 1998–99**




#### **Youth Tobacco Use**

Grades 6–8			GRADES	s 9–12
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
PA <sup>†</sup>	12.0%	15.8%	27.6%	34.0%
Boys <sup>†</sup>	12.5%	18.2%	26.9%	36.8%
Girls <sup>†</sup>	11.4%	13.1%	28.3%	31.0%

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $^{\dagger}$ Pennsylvania Youth Tobacco Survey, 2001.

#### Disparities Among Adult Population Groups, 2000\*





Pennsylvania is investing 66% of CDC's Best Practices lower estimated recommended funding and 23% of the upper estimated recommended funding. Pennsylvania spent \$338 per capita on smoking-attributable direct medical expenditures. In 1998, about 16% (\$1,335,000,000 or \$876.33 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

# Smoking-Attributable Direct Medical Expenditures, 1998

# Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$3,941,000,000 \$329

# State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$341,834,697 Gross cigarette tax revenue collected in 2000 \$333,665,000

Cigarette tax per pack was \$0.31 in 2001

Cigarette sales were 87.9 packs per capita in 2000


FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE	
State Appropriation—Settlement (Tobacco Only)	\$41,402,000	7/01–6/02	
State Appropriation—Excise Tax Revenue	\$0		
State Appropriation—Other	\$0		
Subtotal: State Appropriation	\$41,402,000		
Federal—CDC Office on Smoking and Health	\$1,260,000	6/01–5/02	
Federal—SAMHSA	\$300,000	10/01–9/02	
Non-Government Source—American Legacy Foundation	\$0		
Non-Government Source—RWJF/AMA	\$200,000	6/01–5/02	
Subtotal: Federal/National Sources	\$1,760,000		
FY02 Total Investment in Tobacco Control		\$43,162,000	
CDC Best Practices Recommended Annual Total (Lower E	\$65,568,000		
CDC Best Practices Recommended Annual Total (Upper E	\$184,759,000		
FY02 Per Capita Investment in Tobacco Control			




SMOKING-ATTRIBUTABLE DEATHS, 1999		SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
Overall Men Women Death Rate	1,704 942 763 283.2/100,000	State Lung Cancer All States State CHD* All States	97.2/100,000 90.2/100,000 69.2/100,000 59.7/100,000	Overall Projected Death Rate Projected All	23,875 9,634/100,000
All States	295.5/100,000	State COPD† All States	60.6/100,000 59.7/100,000	Štates	8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

# Adult Cigarette Use, 2000



# **Environmental Tobacco Smoke, 1998-99**




#### **Youth Tobacco Use**

Grades 6–8			GRADES 9–12		
Current Cigarette		Current Any	Current Cigarette	Current Any	
Smoking		Tobacco Use	Smoking	Tobacco Use	
National*	11.0%	15.1%	28.0%	34.5%	
RI <sup>†</sup>	9.1%	13.2%	26.0%	32.1%	
Boys <sup>†</sup>	7.3%	12.4%	26.9%	36.0%	
Girls <sup>†</sup>	10.8%	13.6%	25.2%	27.8%	

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $^{\dagger}$ Rhode Island Youth Tobacco Survey, 2001.

# **Disparities Among Adult Population Groups, 2000\***





Rhode Island is investing 46% of CDC's Best Practices lower estimated recommended funding and 21% of the upper estimated recommended funding. Rhode Island spent \$400 per capita on smoking-attributable direct medical expenditures. In 1998, about 15% (\$140,000,000 or \$912.01 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

# Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory Hospital Nursing Home Prescription Drugs Other Annual Total	\$111,000,000 \$90,000,000 \$119,000,000 \$33,000,000 \$42,000,000 \$395,000,000
Annual Per Capita	\$400

# Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$283,000,000 \$286

# State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$44,589,359 Gross cigarette tax revenue collected in 2000 \$58,788,000

Cigarette tax per pack was \$1.00 in 2001

Cigarette sales were 83.1 packs per capita in 2000


FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$0	
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$3,000,000	7/01–6/02
Subtotal: State Appropriation	\$3,000,000	
Federal—CDC Office on Smoking and Health	\$1,100,000	6/01–5/02
Federal—SAMHSA	\$200,000	10/01–9/02
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$199,819	6/01–5/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		\$4,499,819
CDC Best Practices Recommended Annual Total (Lower E	\$9,888,000	
CDC Best Practices Recommended Annual Total (Upper E	\$21,908,000	
FY02 Per Capita Investment in Tobacco Control		\$4.27




Smoking-Attributable Deaths, 1999		SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
Overall	5,992	State Lung Cancer	96.5/100,000	Overall	90,715
Men Women	3,915 2,077	All States State CHD*	90.2/100,000 64.1/100,000	Projected Death Rate	8,985/100,000
Death Rate All States	315.6/100,000 295.5/100,000	All States State COPD† All States	59.7/100,000 69.5/100,000 59.7/100,000	Projected All States	8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

#### **Adult Cigarette Use, 2000**



# **Environmental Tobacco Smoke, 1998-99**




#### **Youth Tobacco Use**

Grades 6–8			GRADE	s 9–12
Current Cigarette Smoking		Current Any Tobacco Use		
National* SC <sup>†</sup> Boys <sup>†</sup> Girls <sup>†</sup>	11.0% n/a n/a n/a	15.1% n/a n/a n/a	28.0% 36.0% 37.7% 34.2%	34.5% 41.5% 46.4% 36.8%

Current Cigarette Smoking = smoked cigarettes on  $\ge 1$  of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000; †South Carolina Youth Risk Behavior Survey, 1999.

# **Disparities Among Adult Population Groups, 2000\***





South Carolina is investing 14% of CDC's Best Practices lower estimated recommended funding and 5% of the upper estimated recommended funding. South Carolina spent \$223 per capita on smoking-attributable direct medical expenditures. In 1998, about 14% (\$307,000,000 or \$516.11 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

# **Smoking-Attributable** Direct Medical Expenditures, 1998

Ambulatory	\$283,000,000
Hospital	\$238,000,000
Nursing Home	\$182,000,000
Prescription Drugs	\$80,000,000
Other	\$72,000,000
Annual Total	\$855,000,000
Annual Per Capita	\$223

# Smoking-Attributable **Productivity Costs, 1999**

Annual Total Annual Per Capita	\$1,672,000,000 \$430

# State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 Gross cigarette tax revenue collected in 2000 \$72,961,122 \$28,077,000

Cigarette tax per pack was \$0.07 in 2001


Cigarette sales were 103.9 packs per capita in 2000

Funding Source	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$1,620,470	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$62,809	7/01–6/02
Subtotal: State Appropriation	\$1,683,279	
Federal—CDC Office on Smoking and Health	\$1,200,000	6/01–5/02
Federal—SAMHSA	\$190,000	10/01-9/02
Non-Government Source—American Legacy Foundation	\$175,583	10/00–9/01
Non-Government Source—RWJF/AMA	\$0	
Subtotal: Federal/National Sources	\$1,565,583	
FY02 Total Investment in Tobacco Control		\$3,248,862
CDC Best Practices Recommended Annual Total (Lower E	\$23,905,000	
CDC Best Practices Recommended Annual Total (Upper E	\$62,013,000	
FY02 Per Capita Investment in Tobacco Control	\$0.80	


SMOKING-ATTRIBUTABLE DEATHS, 1999		Smoking-Attributable Deaths, 1999 Disease-Specific		s, Youth Projected to Die from Smoking, 1999–2000	
Overall Men Women	1,066 728 338	State Lung Cancer All States State CHD*	70.9/100,000 90.2/100,000 49.1/100,000	Overall Projected Death Rate	20,033 9,886/100,000
Death Rate All States	230.2/100,000 295.5/100,000	All States State COPD† All States	59.7/100,000 54.3/100,000 59.7/100,000	Projected All States	8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

# Adult Cigarette Use, 2000



# **Environmental Tobacco Smoke, 1998–99**




#### **Youth Tobacco Use**

Grades 6–8		GRADES	s <b>9–12</b>	
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
SD	12.4%†	16.3% <sup>†</sup>	32.6% <sup>††</sup>	39.2% <sup>††</sup>
Boys	14.5%†	20.4% <sup>†</sup>	30.4% <sup>††</sup>	41.5% <sup>††</sup>
Girls	10.0%†	11.7% <sup>†</sup>	34.9% <sup>††</sup>	36.9% <sup>††</sup>

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Source: \*National Youth Tobacco Survey, 2000; †South Dakota Youth Tobacco Survey, 1999; ††South Dakota Youth Tobacco Survey, 2000.

# Disparities Among Adult Population Groups, 2000\*





South Dakota is investing 52% of CDC's Best Practices lower estimated recommended funding and 25% of the upper estimated recommended funding. South Dakota spent \$293 per capita on smoking-attributable direct medical expenditures. In 1998, about 13% (\$45,000,000 or \$504.20 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

# Smoking-Attributable Direct Medical Expenditures, 1998

# Ambulatory \$68,000,000 Hospital \$58,000,000 Nursing Home \$63,000,000 Prescription Drugs \$14,000,000 Other \$11,000,000 Annual Total \$214,000,000 Annual Per Capita \$293

# Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$189,000,000 \$258

#### State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$21,642,999 Gross cigarette tax revenue collected in 2000 \$19,251,000

Cigarette tax per pack was \$0.33 in 2001

Cigarette sales were 75.1 packs per capita in 2000


FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$1,000,000	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$2,269,532	7/01–6/02
Subtotal: State Appropriation	\$3,269,532	
Federal—CDC Office on Smoking and Health	\$886,491	6/01–5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$331,533	6/01–5/02
Subtotal: Federal/National Sources	\$1,218,024	
FY02 Total Investment in Tobacco Control		\$4,487,556
CDC Best Practices Recommended Annual Total (Lower E	\$8,688,000	
CDC Best Practices Recommended Annual Total (Upper E	\$18,214,000	
FY02 Per Capita Investment in Tobacco Control	\$5.90	




Smoking-Attributable Deaths, 1999		Smoking-Attributable Deaths, 1999 Disease-Specific		rs, Youth Projected to Die from Smoking, 1999–2000	
Overall Men Women Death Rate	9,624 6,099 3,525 346.0/100,000	State Lung Cancer All States State CHD* All States	110.3/100,000 90.2/100,000 77.3/100,000 59.7/100,000	Overall Projected Death Rate Projected All	124,327 8,890/100,000
All States	295.5/100,000	State COPD† All States	76.9/100,000 59.7/100,000	Štates	8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

# Adult Cigarette Use, 2000



# **Environmental Tobacco Smoke, 1998–99**




#### **Youth Tobacco Use**

GRADES 6–8		GRADES	s 9–12	
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
TN <sup>†</sup>	16.6%	23.2%	32.4%	41.3%
Boys <sup>†</sup>	17.5%	27.0%	33.4%	47.1%
Girls <sup>†</sup>	15.6%	19.1%	31.3%	35.1%

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $\dagger$ Tennessee Youth Tobacco Survey, 2000.

# **Disparities Among Adult Population Groups, 2000\***





Tennessee is investing 6% of CDC's Best Practices lower estimated recommended funding and 2% of the upper estimated recommended funding. Tennessee spent \$311 per capita on smoking-attributable direct medical expenditures. In 1998, about 15% (\$531,000,000 or \$288.17 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

# Smoking-Attributable Direct Medical Expenditures, 1998

# Ambulatory \$615,000,000 Hospital \$406,000,000 Nursing Home \$435,000,000 Prescription Drugs \$139,000,000 Other \$96,000,000 Annual Total \$1,691,000,000 Annual Per Capita \$311

# Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$2,440,000,000 \$445

#### State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001

\$151,394,114 \$79,981,000

Gross cigarette tax revenue collected in 2000

Cigarette tax per pack was \$0.13 in 2001

Cigarette sales were 108.7 packs per capita in 2000


Funding Source	FY02 Amount	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$0	
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$0	
Federal—CDC Office on Smoking and Health	\$1,439,207	6/01–5/02
Federal—SAMHSA	\$125,469	10/01–9/02
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$312,060	6/01–5/02
Subtotal: Federal/National Sources	\$1,876,736	
FY02 Total Investment in Tobacco Control		\$1,876,736
CDC Best Practices Recommended Annual Total (Lower E	\$32,233,000	
CDC Best Practices Recommended Annual Total (Upper E	\$89,079,000	
FY02 Per Capita Investment in Tobacco Control	\$0.33	




Smoking-Attributable Deaths, 1999		Smoking-Attributable Deaths, 1999 Disease-Specific		s, Youth Projected to Die from Smoking, 1999–2000	
Overall Men Women Death Rate	24,158 15,105 9,071 288.8/100,000	State Lung Cancer All States State CHD* All States	87.1/100,000 90.2/100,000 64.1/100,000 59.7/100,000	Overall Projected Death Rate Projected All	486,631 8,267/100,000
All States	295.5/100,000	State COPD† All States	69.9/100,000 59.7/100,000	States	8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

# Adult Cigarette Use, 2000



# **Environmental Tobacco Smoke, 1998-99**




#### **Youth Tobacco Use**

GRADES 6–8			GRADES 9–12	
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
TX <sup>†</sup>	13.9%	17.6%	28.1%	34.6%
Boys <sup>†</sup>	14.6%	19.8%	32.0%	41.8%
Girls <sup>†</sup>	13.2%	15.6%	23.6%	26.3%

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000; †Texas Youth Tobacco Survey, 2000.

# **Disparities Among Adult Population Groups, 2000\***





Texas is investing 16% of CDC's *Best Practices* lower estimated recommended funding and 6% of the upper estimated recommended funding. Texas spent \$231 per capita on smoking-attributable direct medical expenditures. In 1998, about 15% (\$1,265,000,000 or \$543.87 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

# Smoking-Attributable Direct Medical Expenditures, 1998

# Ambulatory \$1,778,000,000 Hospital \$1,046,000,000 Nursing Home \$910,000,000 Prescription Drugs \$397,000,000 Other \$421,000,000 Annual Total \$4,552,000,000 Annual Per Capita \$231

# Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$5,540,000,000 \$276

#### State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$974,220,834 Gross cigarette tax revenue collected in 2000 \$536,877,000

Cigarette tax per pack was \$0.41 in 2001


Cigarette sales were 69.3 packs per capita in 2000

FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$12,500,000	9/01–8/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$1,207,016	9/01–8/02
Subtotal: State Appropriation	\$13,707,016	
Federal—CDC Office on Smoking and Health	\$969,828	6/01–5/02
Federal—SAMHSA	\$1,560,117	10/01–9/02
Non-Government Source—American Legacy Foundation	\$172,500	10/00–9/01
Non-Government Source—RWJF/AMA	\$337,060	6/01–5/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		\$16,746,521
CDC Best Practices Recommended Annual Total (Lower F	\$103,288,000	
CDC Best Practices Recommended Annual Total (Upper E	\$284,735,000	
FY02 Per Capita Investment in Tobacco Control	<b>\$0.79</b>	


SMOKING-ATTRIBUTABLE DEATHS, 1999		SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
Overall Men Women Death Rate	1,218 828 390 159.8/100,000	State Lung Cancer All States State CHD* All States	38.7/100,000 90.2/100,000 27.2/100,000 59.7/100,000	Overall Projected Death Rate Projected All	38,325 5,333/100,000
All States	295.5/100,000	State COPD† All States	50.4/100,000 59.7/100,000	States	8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

# Adult Cigarette Use, 2000



# **Environmental Tobacco Smoke, 1998-99**




#### **Youth Tobacco Use**

GRADES 6–8			GRADES 9–12	
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
UT <sup>†</sup>	n/a	n/a	11.9%	14.5%
Boys <sup>†</sup>	n/a	n/a	11.7%	16.2%
Girls <sup>†</sup>	n/a	n/a	11.8%	12.2%

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $\dagger$ Utah Youth Risk Behavior Survey, 1999.

# **Disparities Among Adult Population Groups, 2000\***





Utah is investing 11% of CDC's *Best Practices* lower estimated recommended funding and 5% of the upper estimated recommended funding. Utah spent \$130 per capita on smoking-attributable direct medical expenditures. In 1998, about 12% (\$81,000,000 or \$373.05 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

# Smoking-Attributable Direct Medical Expenditures, 1998

# Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$244,000,000 \$115

#### **State Revenue from Tobacco Sales and Settlement**

Tobacco settlement revenue received in 2001 \$27,593,646 Gross cigarette tax revenue collected in 2000 \$46,235,000

Cigarette tax per pack was \$0.515 in 2001

Cigarette sales were 40.7 packs per capita in 2000

FUNDING SOURCE	FY02 Amount	Funding Cycle
State Appropriation—Settlement (Tobacco Only)*	\$0	
State Appropriation—Excise Tax Revenue	\$250,000	7/01–6/02
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$250,000	
Federal—CDC Office on Smoking and Health	\$1,268,944	6/01-5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation <sup>†</sup>	\$0	
Non-Government Source—RWJF/AMA	\$199,885	4/01–3/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control	\$1,718,829	
CDC Best Practices Recommended Annual Total (Lower Es	\$15,230,000	
CDC Best Practices Recommended Annual Total (Upper Es	\$33,383,000	
FY02 Per Capita Investment in Tobacco Control	\$0.75	


<sup>\*</sup>A total of \$4,039,800 was appropriated to the Department of Health Community and Family Services for tobacco, alcohol, and drug prevention programs. An additional \$2,000,000 may be appropriated as a supplemental for FY02 per Senate Bill 15 in the upcoming 2002 general session.

†On October 1, 2001, Utah received \$467,635 from the American Legacy Foundation.


SMOKING-ATTRIBUTABLE DEATHS, 1999	SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
Overall         824           Men         517           Women         309           Death Rate         274.3/100,000           All States         295.5/100,000	All States State CHD* All States	90.2/100,000 90.2/100,000 49.4/100,000 59.7/100,000 75.6/100,000 59.7/100,000	Overall Projected Deatl Rate Projected All States	15,801 1 10,711/100,000 8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

# Adult Cigarette Use, 2000



#### **Environmental Tobacco Smoke, 1998-99**




#### **Youth Tobacco Use**

GRADES 6–8			Grades 9–12	
Current C		Current Any	Current Cigarette	Current Any
Smol		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
VT	11.9%†	15.3% <sup>†</sup>	33.4%††	38.6%††
Boys	9.6%†	14.8% <sup>†</sup>	33.3%††	42.1%††
Girls	14.2%†	15.7% <sup>†</sup>	33.6%††	34.8%††

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000; †Vermont Youth Tobacco Survey, 2000; †Vermont Youth Risk Behavior Survey, 1999.

# **Disparities Among Adult Population Groups, 2000\***





Vermont is investing 98% of CDC's Best Practices lower estimated recommended funding and 49% of the upper estimated recommended funding. Vermont spent \$308 per capita on smoking-attributable direct medical expenditures. In 1998, about 14% (\$56,000,000 or \$451.52 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

# Smoking-Attributable Direct Medical Expenditures, 1998

# Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$172,000,000 \$290

# State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$24,458,081 Gross cigarette tax revenue collected in 2000 \$24,144,000

Cigarette tax per pack was \$0.44 in 2001

Cigarette sales were 88.9 packs per capita in 2000

Funding Source	FY02 AMOUNT	FUNDING CYCLE		
State Appropriation—Settlement (Tobacco Only)*	\$5,824,000	7/01–6/02		
State Appropriation—Excise Tax Revenue	\$0			
State Appropriation—Other	\$0			
Subtotal: State Appropriation	\$5,824,000			
Federal—CDC Office on Smoking and Health	\$1,146,500	6/01–5/02		
Federal—SAMHSA	\$0			
Non-Government Source—American Legacy Foundation	\$500,000	10/00–9/01		
Non-Government Source—RWJF/AMA	\$301,584	3/01–2/02		
Subtotal: Federal/National Sources	\$1,948,084			
FY02 Total Investment in Tobacco Control	FY02 Total Investment in Tobacco Control			
CDC Best Practices Recommended Annual Total (Lower 1	\$7,905,000			
CDC Best Practices Recommended Annual Total (Upper 1	\$15,938,000			
FY02 Per Capita Investment in Tobacco Control	\$12.67			


<sup>\*</sup>An additional \$1,028,500 was appropriated for to bacco and other health programs.




SMOKING-ATTRIBUTABLE DEATHS, 1999		SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
Overall Men Women Death Rate All States	9,157 5,604 3,553 290.8/100,000 295.5/100.000	State Lung Cancer All States State CHD* All States State COPD†	94.9/100,000 90.2/100,000 54.0/100,000 59.7/100,000 65.6/100,000	Overall Projected Death Rate Projected All States	134,529 7,739/100,000 8,830/100,000
All States	295.5/100,000	All States	59.7/100,000	States	8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

# Adult Cigarette Use, 2000



# **Environmental Tobacco Smoke, 1998–99**




#### **Youth Tobacco Use**

Grades 6–8			GRADES 9–12		
Current C		Current Any	Current Cigarette	Current Any	
Smok		Tobacco Use	Smoking	Tobacco Use	
National*	11.0%	15.1%	28.0%	34.5%	
VA	n/a	n/a	n/a	n/a	
Boys	n/a	n/a	n/a	n/a	
Girls	n/a	n/a	n/a	n/a	

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Source: \*National Youth Tobacco Survey, 2000.

# **Disparities Among Adult Population Groups, 2000\***





Virginia is investing 44% of CDC's *Best Practices* lower estimated recommended funding and 16% of the upper estimated recommended funding. Virginia spent \$240 per capita on smoking-attributable direct medical expenditures. In 1998, about 14% (\$313,000,000 or \$479.69 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

# Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory	\$597,000,000
Hospital	\$406,000,000
Nursing Home	\$353,000,000
Prescription Drugs	\$150,000,000
Other	\$123,000,000
Annual Total	\$1,629,000,000
Annual Per Capita	\$240

# Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$2,080,000,000 \$303

# State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$126,823,184 Gross cigarette tax revenue collected in 2000 \$16,918,000

Cigarette tax per pack was \$0.025 in 2001

Cigarette sales were 96.7 packs per capita in 2000


FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$15,807,984	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$15,807,984	
Federal—CDC Office on Smoking and Health	\$1,131,145	6/01–5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$0	
Subtotal: Federal/National Sources	\$1,131,145	
FY02 Total Investment in Tobacco Control		\$16,939,129
CDC Best Practices Recommended Annual Total (Lower E	\$38,866,000	
CDC Best Practices Recommended Annual Total (Upper E	\$106,854,000	
FY02 Per Capita Investment in Tobacco Control	\$2.36	




SMOKING-ATTRIBUTABLE DEATHS, 1999		Smoking-Attributable Deaths, 1999 Disease-Specific		Youth Projected to Die from Smoking, 1999–2000	
Overall Men Women Death Rate All States	7,761 4,689 3,072 284.9/100,000 295.5/100,000	State Lung Cancer All States State CHD* All States State COPD† All States	90.1/100,000 90.2/100,000 52.3/100,000 59.7/100,000 76.2/100,000 59.7/100,000	Overall Projected Death Rate Projected All States	133,952 8,848/100,000 8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

#### **Adult Cigarette Use, 2000**



# **Environmental Tobacco Smoke, 1998–99**




#### **Youth Tobacco Use**

Grades 6–8			GRADES 9–12		
Current C		Current Any	Current Cigarette	Current Any	
Smok		Tobacco Use	Smoking	Tobacco Use	
National*	11.0%	15.1%	28.0%	34.5%	
WA	n/a	n/a	n/a	n/a	
Boys	n/a	n/a	n/a	n/a	
Girls	n/a	n/a	n/a	n/a	

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Source: \*National Youth Tobacco Survey, 2000.

# **Disparities Among Adult Population Groups, 2000\***





Washington is investing 62% of CDC's Best Practices lower estimated recommended funding and 23% of the upper estimated recommended funding. Washington spent \$269 per capita on smoking-attributable direct medical expenditures. In 1998, about 16% (\$508,000,000 or \$359.58 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

# Smoking-Attributable Direct Medical Expenditures, 1998

# Smoking-Attributable Productivity Costs, 1999

Annual Total	\$1,503,000,000
Annual Per Capita	\$261

# State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$127,351,202 Gross cigarette tax revenue collected in 2000 \$254,770,000

Cigarette tax per pack was \$0.825 in 2001 and will increase to \$1.425 effective July 2002

Cigarette sales were 52.8 packs per capita in 2000


FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$17,496,000	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$898,500	7/01–6/02
Subtotal: State Appropriation	\$18,394,500	
Federal—CDC Office on Smoking and Health	\$1,524,995	6/01–5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$722,795	10/00–9/01
Non-Government Source—RWJF/AMA	\$0	
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control		\$20,642,290
CDC Best Practices Recommended Annual Total (Lower E	\$33,341,000	
CDC Best Practices Recommended Annual Total (Upper E	\$89,381,000	
FY02 Per Capita Investment in Tobacco Control	\$3.44	




SMOKING-ATTRIBUTABLE DEATHS, 1999		SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		Youth Projected to Die from Smoking, 1999–2000	
Overall Men Women Death Rate	3,838 2,348 1,492 354.1/100,000	State Lung Cancer All States State CHD* All States	90.2/100,000 77.4/100,000 59.7/100,000	Overall Projected Death Rate Projected All	10,571/100,000
All States	295.5/100,000	State COPD† All States	86.7/100,000 59.7/100,000	States	8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

# **Adult Cigarette Use, 2000**



# **Environmental Tobacco Smoke, 1998-99**




#### **Youth Tobacco Use**

GRADES 6–8			GRADES 9–12	
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
WV <sup>†</sup>	18.1%	25.3%	38.5%	47.9%
Boys <sup>†</sup>	17.5%	28.8%	36.0%	52.2%
Girls <sup>†</sup>	18.8%	21.4%	40.6%	42.5%

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $\dagger$ West Virginia Youth Tobacco Survey, 2000.

# **Disparities Among Adult Population Groups, 2000\***





West Virginia is investing 56% of CDC's *Best Practices* lower estimated recommended funding and 23% of the upper estimated recommended funding. West Virginia spent \$298 per capita on smoking-attributable direct medical expenditures. In 1998, about 14% (\$179,000,000 or \$520.55 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

# Smoking-Attributable Direct Medical Expenditures, 1998

# Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$850,000,000 \$470

#### **State Revenue from Tobacco Sales and Settlement**

Tobacco settlement revenue received in 2001 \$54,981,783 Gross cigarette tax revenue collected in 2000 \$33,766,000

Cigarette tax per pack was \$0.17 in 2001

Cigarette sales were 107.9 packs per capita in 2000

FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)*	\$5,850,592	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$5,850,592	
Federal—CDC Office on Smoking and Health	\$1,145,612	6/01–5/02
Federal—SAMHSA	\$35,000	10/01-9/02
Non-Government Source—American Legacy Foundation	\$955,689	10/00–9/01
Non-Government Source—RWJF/AMA	\$0	
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control	\$7,986,893	
CDC Best Practices Recommended Annual Total (Lower E	\$14,160,000	
CDC Best Practices Recommended Annual Total (Upper E	\$35,365,000	
FY02 Per Capita Investment in Tobacco Control	\$4.41	


<sup>\*</sup>An additional \$500,000 was appropriated for the Tobacco/Alcohol Education Program. The appropriation language also states that to the extent permitted by law, eight classified exempt positions shall be provided for the educator-inspector positions to be used in the education and enforcement of activities relating to underage tobacco and alcohol use and sales.




SMOKING-ATTRIBUTABLE DEATHS, 1999		Smoking-Attributable Deaths, 1999 Disease-Specific		Youth Projected to Die from Smoking, 1999–2000	
Overall Men Women Death Rate All States	7,850 4,625 3,225 278.2/100,000 295.5/100,000	State Lung Cancer All States State CHD* All States State COPD† All States	79.6/100,000 90.2/100,000 58.8/100,000 59.7/100,000 61.8/100,000 59.7/100,000	Overall Projected Death Rate Projected All States	136,373 9,963/100,000 8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

# **Adult Cigarette Use, 2000**



#### **Environmental Tobacco Smoke, 1998-99**




#### **Youth Tobacco Use**

	GRADI	ES 6–8	GRADES 9–12	
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
WI <sup>†</sup>	12.2%	16.1%	32.9%	39.4%
Boys <sup>†</sup>	11.9%	17.7%	31.0%	42.1%
Girls <sup>†</sup>	12.7%	14.4%	35.0%	36.2%

Current Cigarette Smoking = smoked cigarettes on  $\geq 1$  of the 30 days preceding the survey. Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on  $\geq 1$  of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000;  $\dagger$ Wisconsin Youth Tobacco Survey, 2000.

# **Disparities Among Adult Population Groups, 2000\***





Wisconsin is investing 24% of CDC's *Best Practices* lower estimated recommended funding and 9% of the upper estimated recommended funding. Wisconsin spent \$303 per capita on smoking-attributable direct medical expenditures. In 1998, about 14% (\$375,000,000 or \$722.80 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

# Smoking-Attributable Direct Medical Expenditures, 1998

# Ambulatory \$570,000,000 Hospital \$322,000,000 Nursing Home \$475,000,000 Prescription Drugs \$121,000,000 Other \$92,000,000 Annual Total \$1,580,000,000 Annual Per Capita \$303

# Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$1,413,000,000 \$269

# State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$123,248,769 Gross cigarette tax revenue collected in 2000 \$251,616,000

Cigarette tax per pack was \$0.77 in 2001

Cigarette sales were 80.1 packs per capita in 2000


FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)*	\$5,532,300	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$5,532,300	
Federal—CDC Office on Smoking and Health	\$1,298,146	6/01–5/02
Federal—SAMHSA	\$76,000	10/01-9/02
Non-Government Source—American Legacy Foundation	\$77,330	10/00–9/01
Non-Government Source—RWJF/AMA	\$485,819	3/01–2/02
Subtotal: Federal/National Sources		
FY02 Total Investment in Tobacco Control	\$7,469,595	
CDC Best Practices Recommended Annual Total (Lower E	\$31,158,000	
CDC Best Practices Recommended Annual Total (Upper E	\$82,381,000	
FY02 Per Capita Investment in Tobacco Control	\$1.38	

<sup>\*</sup>In FY02, \$500,000 was appropriated to the Medical College of Wisconsin for tobacco research. An additional \$9,304,000 was carried over from FY01 to be used for tobacco control in FY02.


SMOKING-ATTRIBUTABLE DEATHS, 1999	SMOKING-ATTRIBUTABLE DEATHS, 1999 DISEASE-SPECIFIC		YOUTH PROJECTED TO DIE FROM SMOKING, 1999–2000	
Overall     729       Men     434       Women     295       Death Rate     318.6/100,000       All States     295.5/100,000	State Lung Cancer All States State CHD* All States State COPD† All States	82.2/100,000 90.2/100,000 50.3/100,000 59.7/100,000 114.2/100,000 59.7/100,000	Overall Projected Deatl Rate Projected All States	12,974 1 10,067/100,000 8,830/100,000

<sup>\*</sup>Coronary heart disease; †chronic obstructive pulmonary disease.

# Adult Cigarette Use, 2000



#### **Environmental Tobacco Smoke, 1998-99**




#### **Youth Tobacco Use**

Grades 6–8			GRADES 9–12	
Current C		Current Any	Current Cigarette	Current Any
Smok		Tobacco Use	Smoking	Tobacco Use
National*	11.0%	15.1%	28.0%	34.5%
WY	14.8% <sup>†</sup>	21.7% <sup>†</sup>	35.2% <sup>††</sup>	43.9%††
Boys	15.6% <sup>†</sup>	25.8% <sup>†</sup>	34.6% <sup>††</sup>	49.6%††
Girls	13.4% <sup>†</sup>	16.9% <sup>†</sup>	35.9% <sup>††</sup>	38.0%††

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥ 1 of the 30 days preceding the survey. Sources: \*National Youth Tobacco Survey, 2000; †Wyoming Youth Tobacco Survey, 2000; †Wyoming Youth Tobacco Survey, 1999.

# **Disparities Among Adult Population Groups, 2000\***





Wyoming is investing 28% of CDC's *Best Practices* lower estimated recommended funding and 14% of the upper estimated recommended funding. Wyoming spent \$221 per capita on smoking-attributable direct medical expenditures. In 1998, about 14% (\$29,000,000 or \$626.16 per recipient) of all Medicaid expenditures were spent on smoking-related illnesses and diseases.

# Smoking-Attributable Direct Medical Expenditures, 1998

Ambulatory	\$33,000,000
Hospital	\$29,000,000
Nursing Home	\$26,000,000
Prescription Drugs	\$10,000,000
Other	\$8,000,000
Annual Total	\$106,000,000
Annual Per Capita	\$221

# Smoking-Attributable Productivity Costs, 1999

Annual Total Annual Per Capita	\$136,000,000 \$284

# State Revenue from Tobacco Sales and Settlement

Tobacco settlement revenue received in 2001 \$14,772,038 Gross cigarette tax revenue collected in 2000 \$5,704,000

Cigarette tax per pack was \$0.12 in 2001

Cigarette sales were 90.5 packs per capita in 2000

FUNDING SOURCE	FY02 AMOUNT	FUNDING CYCLE
State Appropriation—Settlement (Tobacco Only)	\$900,000	7/01–6/02
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$900,000	
Federal—CDC Office on Smoking and Health	\$973,302	6/01–5/02
Federal—SAMHSA	\$0	
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	\$200,000	6/01–5/02
Subtotal: Federal/National Sources	\$1,173,302	
FY02 Total Investment in Tobacco Control		\$2,073,302
CDC Best Practices Recommended Annual Total (Lower Estimate) CDC Best Practices Recommended Annual Total (Upper Estimate)		\$7,381,000
		\$14,397,000
FY02 Per Capita Investment in Tobacco Control		\$4.16